

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



127p  
F0167

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/82, C12Q 1/68, G01N 33/53, A61K 38/17, 48/00, A01K 67/027</b>		A1	(11) International Publication Number: <b>WO 97/19110</b> (43) International Publication Date: 29 May 1997 (29.05.97)
(21) International Application Number: PCT/GB96/02904 (22) International Filing Date: 25 November 1996 (25.11.96) (30) Priority Data: 9523959.6 23 November 1995 (23.11.95) GB 9525555.0 14 December 1995 (14.12.95) GB 9617961.9 28 August 1996 (28.08.96) GB (71) Applicants (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB). DUKE UNIVERSITY [US/US]; Durham, NC 27708 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FUTREAL, Phillip, Andrew [US/US]; Duke University Medical Center, P.O. Box 26111, Durham, NC 27710 (US). WOOSTER, Richard, Francis [GB/GB]; 1 Rose Cottages, Chequiers Lane, Tadworth, Surrey KT20 7ST (GB). ASHWORTH, Alan [GB/GB]; 88B Drayton Gardens, South Kensington SW10 9RF (GB). STRATTON, Michael, Rudolf [GB/GB]; 6 Creighton Road, London NW6 4ED (GB).			(74) Agents: KIDDLE, Simon, J. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: MATERIALS AND METHODS RELATING TO THE IDENTIFICATION AND SEQUENCING OF THE BRCA2 CANCER SUSCEPTIBILITY GENE AND USES THEREOF			
(57) Abstract <p>The identification and sequencing of the BRCA2 gene is disclosed as well as the amino acid sequence of the corresponding BRCA2 polypeptides. BRCA2 alleles including those with mutations in the BRCA2 gene which are associated with a predisposition to develop cancer, especially breast and ovarian cancer, are also disclosed. The present invention further relates to polypeptides encoded by the above nucleic acid. The present invention further relates to uses of such BRCA2 nucleic acid and BRCA2 polypeptides, in particular in the diagnostic, prognostic or therapeutic treatment of cancer.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Materials and Methods Relating to the Identification and Sequencing  
of the BRCA2 Cancer Susceptibility Gene and Uses Thereof

Field of the Invention

5 The present invention relates to the identification and sequencing of  
the BRCA2 cancer susceptibility gene, and to materials and methods  
deriving from these findings. In particular, the present invention  
relates to nucleic acid molecules encoding BRCA2 polypeptides, and  
10 alleles of the BRCA2 gene, including those with mutations which are  
associated with a predisposition to develop cancer, especially breast  
and ovarian cancer, and to polypeptides encoded by this nucleic acid.  
The present invention further relates to uses of such BRCA2 nucleic  
acid and BRCA2 polypeptides, in particular in the diagnostic,  
15 prognostic or therapeutic treatment of cancer.

Background of the Invention

20 Over a lifetime approximately 1 in 12 women develops cancer of the  
breast. While a large majority of these cancers are thought to be  
sporadic, a proportion of breast cancer cases, often quoted at  
approximately 5%, is attributable to a predisposition to the disease  
which is transmitted as a highly penetrant autosomal dominant trait.  
This usually manifests as familial clustering of early onset breast  
25 cancer cases which is often associated with cancers of other organs,  
notably the ovary.

Abnormalities of several genes are known to confer susceptibility to  
breast cancer. The BRCA1 gene is located on chromosome 17q21 (1).  
30 BRCA1 encodes a 1863 aa protein which contains a RING finger domain and  
has little other homology to previously characterised proteins (3).  
Germline mutations of BRCA1 usually result in truncation or absence of  
the protein and hence presumed inactivation of one or more of its  
critical functions. BRCA1 accounts for approximately a third of  
35 families with site specific breast cancer. A small proportion of  
familial breast cancers are attributable to germline mutations in the  
p53 gene and rare clusters of male breast cancers (4) are due to  
mutations in the androgen receptor. Work relating to the BRCA1 gene,  
methods used to isolate it and applications of the BRCA1 nucleic acid  
40 and polypeptides are disclosed in EP-A-0705902.

Using families with multiple cases of early onset breast cancer showing  
evidence against linkage to BRCA1, the present inventors recently  
demonstrated the existence of a second major breast cancer  
45 susceptibility locus, BRCA2, on chromosome 13q12-q13 (5). Preliminary  
studies indicate that mutations in BRCA2 confer a similar risk of  
breast cancer to BRCA1. However, the risk of ovarian cancer appears to

be lower and the risk of male breast cancer substantially higher. Together BRCA1 and BRCA2 account for three quarters of families with multiple early onset breast cancer cases, and almost all families with both breast and ovarian cancer.

5

#### Summary of the Invention

Broadly, the work described in this application is based on the identification of the BRCA2 gene, and the disclosure of its sequence and the amino acid sequence of the corresponding BRCA2 polypeptides. BRCA2 alleles, including those with mutations in the normal sequence, are also disclosed.

The inventors initially found a portion of exon 16 of the BRCA2 gene and used this to sequence approximately 10% of the coding sequence. The BRCA2 cDNA sequence corresponding to this portion of the BRCA2 gene is shown in figure 1, the genomic sequence of the introns and exons initially sequenced by the inventors is shown in figure 2, with the translated protein sequence being shown in figure 3.

20

Just after the identification of 10% of the coding sequence of the BRCA2 gene by the present inventors, a 900,000bp sequence was released on the Internet on 23/11/95 and can be accessed at <ftp://ftp.sanger.ac.uk/pub/human/sequences/13q> and on <ftp://genome.wustl.edu/pub/gsc1/brca2>.

25

Further work by the present inventors using this sequence and other known methods isolated around 75% of the BRCA2 coding sequence. This is shown in the cDNA sequence of figure 4, with the corresponding translated amino acid sequence shown in figure 5. The inventors also at this time identified 6 mutations in the BRCA2 gene, most notably a mutation at 6174delT from analysis of pedigrees from Ashkenazi Jewish families from Montreal.

30

The full BRCA2 sequence, showing the intron and exon structure is set out in figure 7, with the promoter region of the BRCA2 locus shown in figure 7. Primers suitable for amplifying the BRCA2 sequence are shown in figure 8 and are also published on 12/3/1996 in Nature Genetics, 12:333-337, 1996.

40

Following the initial sequencing of the BRCA2 gene described above, and using the information contained in figures 1 to 3, the skilled person could readily assemble the full length sequence of the BRCA2 gene included in the Internet sequence using the techniques described in detail below.

45

In a first aspect, the present invention provides a nucleic acid molecule comprising a part of the BRCA2 gene as set out in figures 1 or 2, or alleles thereof.

5 This nucleic acid can be used to obtain the full length sequence of the BRCA2 gene. Accordingly, in a further aspect, the present invention provides a nucleic acid molecule comprising the full length coding sequence or complete BRCA2 gene as obtainable by:

10 (a) using the nucleic acid sequences shown in figures 1 or 2 to construct probes for screening cDNA or genomic libraries, sequencing the positive clones obtained and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained;

15 (b) using the sequences shown in figures 1 or 2 to obtain oligonucleotides for priming BRCA2 nucleic acid fragments, these oligonucleotides being used in conjunction with oligonucleotides designed to prime from a cloning vector, to amplify by PCR nucleic acid fragments in a library that contains fragments of the BRCA2 sequence, sequencing the amplified fragments to obtain the BRCA2 sequence between known parts of the sequence and the cloning vector, and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained; and/or,

20 (c) using rapid amplification of cDNA ends (RACE), by synthesizing cDNAs from a number of different tissue RNAs, the cDNAs being ligated to an oligonucleotide linker, and amplifying by PCR the BRCA2 cDNAs using one primer that primes from the BRCA2 cDNA sequence of figure 1 and a second primer that primes from the oligonucleotide linker, sequencing the amplified nucleic acid and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained.

30 In a further aspect, the present invention provides a nucleic acid molecule comprising a part of the BRCA2 gene as set out in figure 4, or alleles thereof. The nucleic acid of figure 4 can be used in the same way as the nucleic acid set out in figures 1 and 2 to obtain the coding sequence of the full length BRCA2 gene.

35 The sequences set out in figures 1, 2 and 4 are believed to be a rare alternative splice of BRCA2 including nucleic acid at the 3' end of exon 16 coding for an additional 8 amino acids (ALCDVKAT). The sequence in figure 7 shows what is thought to be the normal amino acid sequence of the BRCA2 polypeptide at this position. However, the presence of the alternative splice has no effect on the methodology outlined above for isolating the full length BRCA2 gene using the 10% and 75% sequences initially isolated by the present inventors.

45

In a further aspect, the present invention provides a nucleic acid molecule which has a nucleotide sequence encoding a BRCA2 polypeptide including the amino acid sequence set out in any one of figures 3, 5 or 7.

5

In a further aspect, the present invention provides a nucleic acid molecule which has a nucleotide sequence encoding a polypeptide which is an allele (including mutant alleles) or variant of a BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5 or 7. Where the nucleic acid sequence is a mutant allele sequence, preferably it includes one or more of the exemplary mutations set out in table 1. Preferred mutations from table 1 include the 6174delT and 6503delTT mutations.

15 In a further aspect, the present invention provides a nucleic acid molecule which has a nucleotide sequence encoding a fragment or active portion of a BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5, or 7.

20 In a further aspect, the present invention provides nucleic acid encoding all or a part the BRCA2 promoter region, the nucleic acid sequence of which is set out in figure 6.

25 In further aspects, the present invention includes replicable vectors comprising the above nucleic acid operably linked to control sequences to direct its expression, host cells transformed with these vectors, and methods of producing BRCA2 polypeptide comprising culturing the host cells and recovering the polypeptide produced.

30 In a further aspect, the present invention provides the above nucleic acid molecules for use in methods of medical treatment, especially in the diagnosis and therapy of cancer. Also included herein is the use of the above nucleic acid molecules in the preparation of a medicament for treating cancer. This is discussed further below.

35

In a further aspect, the present invention provides the use of one of the above nucleic acid sequences in the design of primers for use in the polymerase chain reaction.

40 In a further aspect, the present invention provides substances comprising polypeptides encoded by the above nucleic acid, or an active portions, derivatives or functional mimetics thereof.

45 In a further aspect, the present invention provides a method of diagnosing a susceptibility or predisposition to cancer in a patient by

analysing a sample from the patient for the BRCA2 gene or the polypeptide encoded by it. By way of example, this could be carried out by:

5 (a) comparing the sequence of nucleic acid in the sample with the BRCA2 nucleic acid sequence to determine whether the sample from the patient contains mutations; or,

(b) determining the presence in a sample from a patient of the polypeptide encoded by the BRCA2 gene as set out in the partial sequences of figures 3 and 5 or the full length sequence set out in figure 7 and, if present, determining whether the polypeptide is full length, and/or is mutated, and/or is expressed at the normal level; or,

10 (c) using DNA fingerprinting to compare the restriction pattern produced when a restriction enzyme cuts a sample of nucleic acid from the patient with the restriction pattern obtained from normal BRCA2 gene comprising the sequence set out in figures 1, 2, 4 or 7 or from known mutations thereof; or,

(d) using a specific binding member capable of binding to a BRCA2 nucleic acid sequence (either a normal sequence or a known mutated sequence), the specific binding member comprising nucleic acid hybridisable with the BRCA2 sequence, or substances comprising an antibody domain with specificity for a native or mutated BRCA2 nucleic acid sequence or the polypeptide encoded by it, and detecting the binding of the specific binding member to its binding partner by means of a label; or,

25 (e) using PCR involving one or more primers based on normal or mutated BRCA2 gene sequence to screen for normal or mutant BRCA2 gene in a sample from a patient.

30 While diagnostic methods (a)-(e) are provided as examples, other assay formats are well known in the art and will be apparent to the skilled person.

35 The detection of mutations in the BRCA2 gene indicates a susceptibility to cancer, especially female breast cancer, male breast cancer and ovarian cancer. Risks of other cancers including prostate cancer, pancreatic cancer, ocular melanoma, colorectal cancer and leukaemia are also likely to be elevated in carriers of BRCA2 mutations.

#### Brief Description of the Drawings

40 The above and further aspects of the present invention will now be further described by way of example with reference to the accompanying drawings, by way of example and not limitation. Still further aspects of the invention will be apparent to those of ordinary skill in the art.

Figure 1 shows the partial sequence of the BRCA2 gene obtained from a cDNA clone 14.

5 Figure 2(a)-(e) shows the sequences of the exons and introns of clone 14 where they are known.

Figure 3 shows the translated amino acid sequence of nucleic acid sequence shown in figure 1. Of the 3 possible reading frames, the one used to translate the amino acid sequence shown in this figure is the most likely candidate as the nucleic acid obtained when the reading frame is in the other positions contains stop codons in the sequence.

10 Figure 4 shows the second part of the cDNA sequence of the BRCA2 gene isolated by the inventors, including the cDNA sequence set out in figure 1.

Figure 5(a)-(d) shows the translated amino acid sequence of nucleic acid sequence shown in figure 4, with the bold arrows over the sequence indicating the boundaries between the parts of the protein encoded by different exons in the gene.

20 Figure 6 shows the sequences of the BRCA2 promoter region with the binding sites for potential transcription factors underlined and CpG shown in bold. The promoter sequence may be of use in screening for substances which modulate the expression of the BRCA2 gene for use as therapeutics.

Figure 7 shows the sequence of the BRCA2 gene, including the exon/intron structure defined by comparing the BRCA2 cDNA sequence with the genomic sequence of chromosome 13q between D13S260 and D13S171. Exons are shown in upper case with the flanking intron sequence shown in lower case. The amino acid sequence is shown below the open reading frame from exon 2 to 26.

35 Figure 8 shows primers for single stranded conformation polymorphism (SSCP) testing which have been designed to amplify the BRCA2 gene by the PCR for the identification of sequence changes in genomic DNA that may predispose to the development of breast cancer. The primers are located in the intron sequences flanking each of the 26 exons that contain the open reading frame of the gene. The amplified products include the splice site consensus sequences. Two or more sets of primers have been designed for the larger exons of the gene. The primers are labelled by: their exon number, the subsection of the exon and as forward or reverse. The PCR products range from 160 to 360bp.

40

45 The column labelled CON indicates the conditions that we have used.



The primers have been tested using a limited set of conditions which were all based on touchdown (TD) PCR where the figures indicate the first and final annealing temperature respectively. When mutations have been identified, the PCR products have been sequenced using the same primers and fluorescent cycle sequencing.

Figure 9 shows the alignment of amino acid motifs in exon 11 of BRCA2 from different species.

Figure 10 shows exemplary primers for use in PTT assays.

Figure 11 shows immunoprecipitation of FLAG-tagged BRCA2 protein from COS cells transfected with expression plasmid p13120. Total cell lysates from transfected, (lanes a and b) and untransfected (lanes c and d) COS cells subjected to immunoprecipitation with anti-FLAG antibody (lanes b and d) or with a control antibody against a Golgi protein (lanes a and c). Anti-FLAG antibody immunoprecipitates a protein of approximately 400kDa (arrow) from transfected, but not from untransfected COS cell lysate (compare lanes d and b). The control antibody does not immunoprecipitate this protein (see lane c). The strong, high molecular weight protein in the marker lanes is 220kDa.

Figure 12 shows a Western blot of FLAG-tagged BRCA2 protein. Total cell lysates of COS cells (lanes a and b) and 293T cells (lanes c and d) transfected with expression plasmid p13120 (lanes a and c) or no DNA (lanes b and d). Anti-FLAG antibody detects a protein of approximately 400kDa (arrow) in the transfected, but not in the untransfected cell lysates. The protein in the marker lane has a molecular weight of 220kDa.

Figure 13 shows the detection of FLAG-tagged BRCA2 protein by immunofluorescence. COS cells transfected with expression plasmid, p13120 and fixed, 48 hours later, with 4% formaldehyde, then permeabilised with 0.4% Triton-X100. Anti-FLAG antibody reveals variation in the cellular localisation of FLAG-tagged BRCA2 protein. Whilst BRCA2 protein is equally distributed between the nucleus and the cytoplasm in 14% of cells, it is predominantly nuclear (A) in 42% and cytoplasmic (B) in 44% of cells (n=50). Expression in the cytoplasm appears granular, which may indicate an association with the endoplasmic reticulum. Staining is not seen in untransfected cells.

#### Detailed Description

Preparation BRCA2 nucleic acid, and vectors and host cells incorporating the nucleic acid

"BRCA2 region" refers to the portion of human chromosome 13q12-13 identified in Wooster et al (5), containing the BRCA2 locus.

5 The "BRCA2 locus" includes the BRCA2 gene, both the coding sequence (exons) and intervening sequences (introns), and its regulatory elements for controlling transcription and/or translation. The BRCA2 locus covers allelic variations within the locus.

10 The term "BRCA2 gene" or "BRCA2 nucleic acid" includes normal alleles of the BRCA2 gene, both silent alleles having no effect on the amino acid sequence of the BRCA2 polypeptide and alleles leading to amino acid sequence variants of BRCA2 polypeptide that do not substantially affect its function. These terms also includes alleles having one or  
15 more mutations that are linked to a predisposition to develop cancer, especially male and female breast cancer or ovarian cancer. A mutation may be a change in the BRCA2 nucleic acid sequence which produces a deleterious change in the amino acid sequence of the BRCA2 polypeptide, resulting in partial or complete loss of BRCA2 function, or may be a change in the nucleic acid sequence which results in the loss of  
20 effective BRCA2 expression or the production of aberrant forms of the BRCA2 polypeptide. Examples of such mutations are shown in table 1. Alleles including such mutations are also known in the art as susceptibility alleles.

25 The BRCA2 nucleic acid may be that shown in figures 1, 2, 4 or 7, or it may be an allele as described above, or a variant or derivative, differing from that shown by a change which is one or more of addition, insertion, deletion and substitution of one or more nucleotides of the sequence shown. Changes to a nucleotide sequence may result in an  
30 amino acid change at the protein level, or not, as determined by the genetic code.

35 Thus, nucleic acid according to the present invention may include a sequence different from the sequence shown in figures 1, 2, 4 or 7 yet encode a polypeptide with the same amino acid sequence. The amino acid sequence shown in figure 7 consists of 3418 residues.

40 On the other hand, the encoded polypeptide may comprise an amino acid sequence which differs by one or more amino acid residues from the amino acid sequence shown in figures 3, 5 or 7. Nucleic acid encoding a polypeptide which is an amino acid sequence variant, derivative or allele of the sequence shown in figures 3, 5 or 7 is further provided by the present invention. Such polypeptides are discussed below. Nucleic acid encoding such a polypeptide may show greater than about  
45 60% homology with the coding sequence shown in figures 1, 2, 4 or 7

greater than about 70% homology, greater than about 80% homology, greater than about 90% homology or greater than about 95% homology.

5 Particular mutant alleles of the present invention are set out in table 1, using the nomenclature first proposed in (8). These mutations are generally associated with the production of truncated forms of the BRCA2 gene product. These have been shown by the experimental work described herein to be associated with susceptibility to male and female breast cancer and/or ovarian cancer. Implications for  
10 screening, e.g. for diagnostic or prognostic purposes, are discussed below.

Generally, nucleic acid according to the present invention is provided as an isolate, in isolated and/or purified form, or free or  
15 substantially free of material with which it is naturally associated, such as free or substantially free of nucleic acid flanking the gene in the human genome, except possibly one or more regulatory sequence(s) for expression. Nucleic acid may be wholly or partially synthetic and may include genomic DNA, cDNA or RNA. Where nucleic acid according to  
20 the invention includes RNA, reference to the sequence shown should be construed as reference to the RNA equivalent, with U substituted for T.

Nucleic acid sequences encoding all or part of the BRCA2 gene and/or its regulatory elements can be readily prepared by the skilled person  
25 using the information and references contained herein and techniques known in the art (for example, see Sambrook, Fritsch and Maniatis, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press, 1989, and Ausubel et al, Short Protocols in Molecular Biology, John Wiley and Sons, 1992). These techniques include (i) the use of  
30 the polymerase chain reaction (PCR) to amplify samples of such nucleic acid, e.g. from genomic sources, (ii) chemical synthesis, or (iii) preparing cDNA sequences. Modifications to the BRCA2 sequences can be made, e.g. using site directed mutagenesis, to provide expression of modified BRCA2 polypeptide or to take account of codon preference in  
35 the host cells used to express the nucleic acid.

In order to obtain expression of the BRCA2 nucleic acid sequences, the sequences can be incorporated in a vector having control sequences operably linked to the BRCA2 nucleic acid to control its expression.  
40 The vectors may include other sequences such as promoters or enhancers to drive the expression of the inserted nucleic acid, nucleic acid sequences so that the BRCA2 polypeptide is produced as a fusion and/or nucleic acid encoding secretion signals so that the polypeptide produced in the host cell is secreted from the cell. BRCA2 polypeptide  
45 can then be obtained by transforming the vectors into host cells in

which the vector is functional, culturing the host cells so that the BRCA2 polypeptide is produced and recovering the BRCA2 polypeptide from the host cells or the surrounding medium. Prokaryotic and eukaryotic cells are used for this purpose in the art, including strains of E. coli, yeast, and eukaryotic cells such as COS or CHO cells. The choice of host cell can be used to control the properties of the BRCA2 polypeptide expressed in those cells, e.g. controlling where the polypeptide is deposited in the host cells or affecting properties such as its glycosylation.

PCR techniques for the amplification of nucleic acid are described in US Patent No. 4,683,195. In general, such techniques require that sequence information from the ends of the target sequence is known to allow suitable forward and reverse oligonucleotide primers to be designed to be identical or similar to the polynucleotide sequence that is the target for the amplification. PCR comprises steps of denaturation of template nucleic acid (if double-stranded), annealing of primer to target, and polymerisation. The nucleic acid probed or used as template in the amplification reaction may be genomic DNA, cDNA or RNA. PCR can be used to amplify specific sequences from genomic DNA, specific RNA sequences and cDNA transcribed from mRNA, bacteriophage or plasmid sequences. The BRCA2 nucleic acid sequences provided herein readily allow the skilled person to design PCR primers, see for example figure 8. References for the general use of PCR techniques include Mullis et al, Cold Spring Harbor Symp. Quant. Biol., 51:263, (1987), Ehrlich (ed), PCR technology, Stockton Press, NY, 1989, Ehrlich et al, Science, 252:1643-1650, (1991), PCR protocols; A Guide to Methods and Applications, Eds. Innis et al, Academic Press, New York, (1990).

Also included within the scope of the invention are antisense oligonucleotide sequences based on the BRCA2 nucleic acid sequences described herein. Antisense oligonucleotides may be designed to hybridise to the complementary sequence of nucleic acid, pre-mRNA or mature mRNA, interfering with the production of polypeptide encoded by a given DNA sequence (e.g. either native BRCA2 polypeptide or a mutant form thereof), so that its expression is reduced or prevented altogether. In addition to the BRCA2 coding sequence, antisense techniques can be used to target the control sequences of the BRCA2 gene, e.g. in the 5' flanking sequence of the BRCA2 coding sequence, whereby the antisense oligonucleotides can interfere with BRCA2 control sequences. The construction of antisense sequences and their use is described in Peyman and Ulman, Chemical Reviews, 90:543-584, (1990), Crooke, Ann. Rev. Pharmacol. Toxicol., 32:329-376, (1992), and Zamecnik and Stephenson, P.N.A.S., 75:280-284, (1974).

The nucleic acid sequences provided in figures 1, 2, 4 and 7 are useful for identifying nucleic acid of interest (and which may be according to the present invention) in a test sample. The present invention provides a method of obtaining nucleic acid of interest, the method including hybridisation of a probe having the sequence shown in figures 1, 2, 4 or 7 or a complementary sequence, to target nucleic acid.

Hybridisation is generally followed by identification of successful hybridisation and isolation of nucleic acid which has hybridised to the probe, which may involve one or more steps of PCR.

Nucleic acid according to the present invention is obtainable using one or more oligonucleotide probes or primers designed to hybridise with one or more fragments of the nucleic acid sequence shown in figures 1, 2, 4 or 7, particularly fragments of relatively rare sequence, based on codon usage or statistical analysis. A primer designed to hybridise with a fragment of the nucleic acid sequence shown in the above figures may be used in conjunction with one or more oligonucleotides designed to hybridise to a sequence in a cloning vector within which target nucleic acid has been cloned, or in so-called "RACE" (rapid amplification of cDNA ends) in which cDNA's in a library are ligated to an oligonucleotide linker and PCR is performed using a primer which hybridises with the sequence shown in figures 1, 2, 4 or 7 and a primer which hybridises to the oligonucleotide linker.

Such oligonucleotide probes or primers, as well as the full-length sequence (and alleles, variants and derivatives) are also useful in screening a test sample containing nucleic acid for the presence of alleles and variants, especially those that confer susceptibility or predisposition to cancers, the probes hybridising with a target sequence from a sample obtained from the individual being tested. The conditions of the hybridisation can be controlled to minimise non-specific binding, and preferably stringent to moderately stringent hybridisation conditions are preferred. The skilled person is readily able to design such probes, label them and devise suitable conditions for the hybridisation reactions, assisted by textbooks such as Sambrook et al (1989) and Ausubel et al (1992).

As well as determining the presence of polymorphisms or mutations in the BRCA2 sequence, the probes may also be used to determine whether mRNA encoding BRCA2 is present in a cell or tissue.

Nucleic acid isolated and/or purified from one or more cells (e.g. human) or a nucleic acid library derived from nucleic acid isolated and/or purified from cells (e.g. a cDNA library derived from mRNA

isolated from the cells), may be probed under conditions for selective hybridisation and/or subjected to a specific nucleic acid amplification reaction such as the polymerase chain reaction (PCR).

5 In the context of cloning, it may be necessary for one or more gene fragments to be ligated to generate a full-length coding sequence. Also, where a full-length encoding nucleic acid molecule has not been obtained, a smaller molecule representing part of the full molecule, may be used to obtain full-length clones. Inserts may be prepared from  
10 partial cDNA clones and used to screen cDNA libraries. The full-length clones isolated may be subcloned into expression vectors and activity assayed by transfection into suitable host cells, e.g. with a reporter plasmid.

15 A method may include hybridisation of one or more (e.g. two) probes or primers to target nucleic acid. Where the nucleic acid is double-stranded DNA, hybridisation will generally be preceded by denaturation to produce single-stranded DNA. The hybridisation may be as part of a PCR procedure, or as part of a probing procedure not involving PCR. An  
20 example procedure would be a combination of PCR and low stringency hybridisation. A screening procedure, chosen from the many available to those skilled in the art, is used to identify successful hybridisation events and isolated hybridised nucleic acid.

25 Binding of a probe to target nucleic acid (e.g. DNA) may be measured using any of a variety of techniques at the disposal of those skilled in the art. For instance, probes may be radioactively, fluorescently or enzymatically labelled. Other methods not employing labelling of probe include examination of restriction fragment length polymorphisms,  
30 amplification using PCR, RNAase cleavage and allele specific oligonucleotide probing.

Probing may employ the standard Southern blotting technique. For instance DNA may be extracted from cells and digested with different  
35 restriction enzymes. Restriction fragments may then be separated by electrophoresis on an agarose gel, before denaturation and transfer to a nitrocellulose filter. Labelled probe may be hybridised to the DNA fragments on the filter and binding determined. DNA for probing may be prepared from RNA preparations from cells.

40 Preliminary experiments may be performed by hybridising under low stringency conditions various probes to Southern blots of DNA digested with restriction enzymes. Suitable conditions would be achieved when a large number of hybridising fragments were obtained while the  
45 background hybridisation was low. Using these conditions nucleic acid

libraries, e.g. cDNA libraries representative of expressed sequences, may be searched.

5 Those skilled in the art are well able to employ suitable conditions of the desired stringency for selective hybridisation, taking into account factors such as oligonucleotide length and base composition, temperature and so on.

10 On the basis of amino acid sequence information, oligonucleotide probes or primers may be designed, taking into account the degeneracy of the genetic code, and, where appropriate, codon usage of the organism from the candidate nucleic acid is derived. An oligonucleotide for use in nucleic acid amplification may have about 10 or fewer codons (e.g. 6, 7 or 8), i.e. be about 30 or fewer nucleotides in length (e.g. 18, 21  
15 or 24). Generally specific primers are upwards of 14 nucleotides in length, but not more than 18-20. Those skilled in the art are well versed in the design of primers for use processes such as PCR.

20 A further aspect of the present invention provides an oligonucleotide or polynucleotide fragment of the nucleotide sequence shown in figure 1, 2, 4 or 7, or a complementary sequence, in particular for use in a method of obtaining and/or screening nucleic acid. The sequences referred to above may be modified by addition, substitution, insertion or deletion of one or more nucleotides, but preferably without  
25 abolition of ability to hybridise selectively with nucleic acid with the sequence shown in figures 1, 2, 4 or 7, that is wherein the degree of homology of the oligonucleotide or polynucleotide with one of the sequences given is sufficiently high.

30 In some preferred embodiments, oligonucleotides according to the present invention that are fragments of any of the sequences shown in figures 1, 2, 4 or 7, or any allele associated with cancer susceptibility, are at least about 10 nucleotides in length, more preferably at least about 15 nucleotides in length, more preferably at  
35 least about 20 nucleotides in length. Such fragments themselves individually represent aspects of the present invention. Fragments and other oligonucleotides may be used as primers or probes as discussed but may also be generated (e.g. by PCR) in methods concerned with determining the presence in a test sample of a sequence indicative of  
40 cancer susceptibility.

Methods involving use of nucleic acid in diagnostic and/or prognostic contexts, for instance in determining susceptibility to cancer, and other methods concerned with determining the presence of sequences  
45 indicative of cancer susceptibility are discussed below.

Nucleic acid according to the present invention may be used in methods of gene therapy, for instance in treatment of individuals with the aim of preventing or curing (wholly or partially) cancer. This too is discussed below.

5 Nucleic acid according to the present invention, such as a full-length coding sequence or oligonucleotide probe or primer, may be provided as part of a kit, e.g. in a suitable container such as a vial in which the contents are protected from the external environment. The kit may  
10 include instructions for use of the nucleic acid, e.g. in PCR and/or a method for determining the presence of nucleic acid of interest in a test sample. A kit wherein the nucleic acid is intended for use in PCR may include one or more other reagents required for the reaction, such as polymerase, nucleosides, buffer solution etc. The nucleic acid may  
15 be labelled. A kit for use in determining the presence or absence of nucleic acid of interest may include one or more articles and/or reagents for performance of the method, such as means for providing the test sample itself, e.g. a swab for removing cells from the buccal cavity or a syringe for removing a blood sample (such components  
20 generally being sterile). In a further aspect, the present invention provides an apparatus for screening BRCA2 nucleic acid, the apparatus comprising storage means including the BRCA2 nucleic acid sequence as set out in any one of figures 1, 2, 4 or 7, the stored sequence being used to compare the sequence of the test nucleic acid to determine the  
25 presence of mutations.

A convenient way of producing a polypeptide according to the present invention is to express nucleic acid encoding it, by use of the nucleic acid in an expression system. The use of expression system has reached  
30 an advanced degree of sophistication today.

Accordingly, the present invention also encompasses a method of making a polypeptide (as disclosed), the method including expression from nucleic acid encoding the polypeptide (generally nucleic acid according  
35 to the invention). This may conveniently be achieved by growing a host cell in culture, containing such a vector, under appropriate conditions which cause or allow expression of the polypeptide. Polypeptides may also be expressed in in vitro systems, such as reticulocyte lysate.

40 Systems for cloning and expression of a polypeptide in a variety of different host cells are well known. Suitable host cells include bacteria, eukaryotic cells such as mammalian and yeast, and baculovirus systems. Mammalian cell lines available in the art for expression of a heterologous polypeptide include Chinese hamster ovary cells, HeLa  
45 cells, baby hamster kidney cells, COS cells and many others. A common,



preferred bacterial host is *E. coli*.

Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. Vectors may be plasmids, viral e.g. 'phage, or phagemid, as appropriate. For further details see, for example, *Molecular Cloning: a Laboratory Manual*: 2nd edition, Sambrook et al., 1989, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in *Current Protocols in Molecular Biology*, Ausubel et al. eds., John Wiley & Sons, 1992.

Thus, a further aspect of the present invention provides a host cell containing nucleic acid as disclosed herein. The nucleic acid of the invention may be integrated into the genome (e.g. chromosome) of the host cell. Integration may be promoted by inclusion of sequences which promote recombination with the genome, in accordance with standard techniques. The nucleic acid may be on an extra-chromosomal vector within the cell.

A still further aspect provides a method which includes introducing the nucleic acid into a host cell. The introduction, which may (particularly for in vitro introduction) be generally referred to without limitation as "transformation", may employ any available technique. For eukaryotic cells, suitable techniques may include calcium phosphate transfection, DEAE-Dextran, electroporation, liposome-mediated transfection and transduction using retrovirus or other virus, e.g. vaccinia or, for insect cells, baculovirus. For bacterial cells, suitable techniques may include calcium chloride transformation, electroporation and transfection using bacteriophage. As an alternative, direct injection of the nucleic acid could be employed.

Marker genes such as antibiotic resistance or sensitivity genes may be used in identifying clones containing nucleic acid of interest, as is well known in the art.

The introduction may be followed by causing or allowing expression from the nucleic acid, e.g. by culturing host cells (which may include cells actually transformed although more likely the cells will be descendants of the transformed cells) under conditions for expression of the gene, so that the encoded polypeptide is produced. If the polypeptide is

expressed coupled to an appropriate signal leader peptide it may be secreted from the cell into the culture medium. Following production by expression, a polypeptide may be isolated and/or purified from the host cell and/or culture medium, as the case may be, and subsequently  
5 used as desired, e.g. in the formulation of a composition which may include one or more additional components, such as a pharmaceutical composition which includes one or more pharmaceutically acceptable excipients, vehicles or carriers (e.g. see below).

10 Introduction of nucleic acid may take place in vivo by way of gene therapy, as discussed below.

A host cell containing nucleic acid according to the present invention, e.g. as a result of introduction of the nucleic acid into the cell or  
15 into an ancestor of the cell and/or genetic alteration of the sequence endogenous to the cell or ancestor (which introduction or alteration may take place in vivo or ex vivo), may be comprised (e.g. in the soma) within an organism which is an animal, particularly a mammal, which may be human or non-human, such as rabbit, guinea pig, rat, mouse or other  
20 rodent, cat, dog, pig, sheep, goat, cattle or horse, or which is a bird, such as a chicken. Genetically modified or transgenic animals or birds comprising such a cell are also provided as further aspects of the present invention.

25 This may have a therapeutic aim. (Gene therapy is discussed below.) The presence of a mutant, allele or variant sequence within cells of an organism, particularly when in place of a homologous endogenous sequence, may allow the organism to be used as a model in testing and/or studying the role of the BRCA2 gene or substances which modulate  
30 activity of the encoded polypeptide in vitro or the promoter sequence shown in figure 6 are otherwise indicated to be of therapeutic potential.

Instead of or as well as being used for the production of a polypeptide  
35 encoded by a transgene, host cells may be used as a nucleic acid factory to replicate the nucleic acid of interest in order to generate large amounts of it. Multiple copies of nucleic acid of interest may be made within a cell when coupled to an amplifiable gene such as DHFR. Host cells transformed with nucleic acid of interest, or which are  
40 descended from host cells into which nucleic acid was introduced, may be cultured under suitable conditions, e.g. in a fermenter, taken from the culture and subjected to processing to purify the nucleic acid. Following purification, the nucleic acid or one or more fragments thereof may be used as desired, for instance in a diagnostic or  
45 prognostic assay as discussed elsewhere herein.

Production of BRCA2 Polypeptides

5 The skilled person can use the techniques described herein and others well known in the art to produce large amounts of the BRCA2 polypeptide, or fragments or active portions thereof, for use as pharmaceuticals, in the developments of drugs and for further study into its properties and role in vivo. Experimental work confirming the production of BRCA2 polypeptide is set out in example 3 below.

10 Thus, a further aspect of the present invention provides a polypeptide which has the amino acid sequence shown in figures 3, 5 or 7, which may be in isolated and/or purified form, free or substantially free of material with which it is naturally associated, such as other polypeptides or such as human polypeptides other than BRCA2 polypeptide  
15 or (for example if produced by expression in a prokaryotic cell) lacking in native glycosylation, e.g. unglycosylated.

Polypeptides which are amino acid sequence variants, alleles or derivatives are also provided by the present invention. A polypeptide  
20 which is a variant, allele, or derivative may have an amino acid sequence which differs from that given in figures 3, 5 or 7 by one or more of addition, substitution, deletion and insertion of one or more amino acids. Preferred such polypeptides have BRCA2 function, that is to say have one or more of the following properties: immunological  
25 cross-reactivity with an antibody reactive the polypeptide for which the sequence is given in figures 3, 5 or 7; sharing an epitope with the polypeptide for which the amino acid sequence is shown in figures 3, 5 or 7 (as determined for example by immunological cross-reactivity between the two polypeptides).

30 A polypeptide which is an amino acid sequence variant, or allele, derivative of the amino acid sequence shown in figures 3, 5, or 7 may comprise an amino acid sequence which shares greater than about 35% sequence identity with the sequence shown, greater than about 40%,  
35 greater than about 50%, greater than about 60%, greater than about 70%, greater than about 80%, greater than about 90% or greater than about 95%. The sequence may share greater than about 60% similarity, greater than about 70% similarity, greater than about 80% similarity or greater than about 90% similarity with the amino acid sequence shown in any one  
40 of figures 3, 5 or 7. Particular amino acid sequence variants may differ from those shown in figures 3, 5 or 7 by insertion, addition, substitution or deletion of 1 amino acid, 2, 3, 4, 5-10, 10-20 20-30, 30-50, 50-100, 100-150, or more than 150 amino acids.

45 By way of example, mutation variants representative of preferred

embodiments of the present invention are shown in table 1. Screening for the presence of one or more of these in a test sample has a diagnostic and/or prognostic use, for instance in determining cancer susceptibility, as discussed below.

5

The present invention also includes active portions, fragments, derivatives and functional mimetics of the BRCA2 polypeptides of the invention.

10

An "active portion" of BRCA2 polypeptide means a peptide which is less than said full length BRCA2 polypeptide, but which retains its essential biological activity.

15

A "fragment" of the BRCA2 polypeptide means a stretch of amino acid residues of at least about five to seven contiguous amino acids, often at least about seven to nine contiguous amino acids, typically at least about nine to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids. Fragments of the BRCA2 polypeptide sequence antigenic determinants or epitopes useful for raising antibodies to a portion of the BRCA2 amino acid sequence.

20

A "derivative" of the BRCA2 polypeptide or a fragment thereof means a polypeptide modified by varying the amino acid sequence of the protein, e.g. by manipulation of the nucleic acid encoding the protein or by altering the protein itself. Such derivatives of the natural amino acid sequence may involve insertion, addition, deletion or substitution of one or more amino acids, without fundamentally altering the essential activity of the wild type BRCA2 polypeptide.

25

30

"Functional mimetic" means a substance which may not contain an active portion of the BRCA2 amino acid sequence, and probably is not a peptide at all, but which retains the essential biological activity of natural BRCA2 polypeptide. The design and screening of candidate mimetics is described in detail below.

35

A polypeptide according to the present invention may be isolated and/or purified (e.g. using an antibody) for instance after production by expression from encoding nucleic acid (for which see below). Polypeptides according to the present invention may also be generated wholly or partly by chemical synthesis. The isolated and/or purified polypeptide may be used in formulation of a composition, which may include at least one additional component, for example a pharmaceutical composition including a pharmaceutically acceptable excipient, vehicle or carrier. A composition including a polypeptide according to the invention may be used in prophylactic and/or therapeutic treatment as

40

45

discussed below.

5 A polypeptide, peptide fragment, allele, or variant according to the present invention may be used as an immunogen or otherwise in obtaining specific antibodies. Antibodies are useful in purification and other manipulation of polypeptides and peptides, diagnostic screening and therapeutic contexts. This is discussed further below.

10 A polypeptide according to the present invention may be used in screening for molecules which affect or modulate its activity or function. Such molecules may be useful in a therapeutic (possibly including prophylactic) context.

#### Production of BRCA2 Antibodies

15 A further important use of the BRCA2 polypeptides is in raising antibodies that have the property of specifically binding to the BRCA2 polypeptides, or fragments or active portions thereof.

20 The production of monoclonal antibodies is well established in the art. Monoclonal antibodies can be subjected to the techniques of recombinant DNA technology to produce other antibodies or chimeric molecules which retain the specificity of the original antibody. Such techniques may involve introducing DNA encoding the immunoglobulin variable region, or  
25 the complementarity determining regions (CDRs), of an antibody to the constant regions, or constant regions plus framework regions, of a different immunoglobulin. See, for instance, EP-A-184187, GB-A-2188638 or EP-A-239400. A hybridoma producing a monoclonal antibody may be subject to genetic mutation or other changes, which may or may not  
30 alter the binding specificity of antibodies produced.

The provision of the novel BRCA2 polypeptides enables for the first time the production of antibodies able to bind it specifically. Accordingly, a further aspect of the present invention provides an  
35 antibody able to bind specifically to the polypeptide whose sequence is given in figures 3, 5 or 7. Such an antibody may be specific in the sense of being able to distinguish between the polypeptide it is able to bind and other human polypeptides for which it has no or substantially no binding affinity (e.g. a binding affinity of about  
40 1000x worse). Specific antibodies bind an epitope on the molecule which is either not present or is not accessible on other molecules. Antibodies according to the present invention may be specific for the wild-type polypeptide. Antibodies according to the invention may be specific for a particular variant, allele or derivative polypeptide as  
45 between that molecule and the wild-type BRCA2 polypeptide, so as to be

useful in diagnostic and prognostic methods as discussed below. Antibodies are also useful in purifying the polypeptide or polypeptides to which they bind, e.g. following production by recombinant expression from encoding nucleic acid.

5

Preferred antibodies according to the invention are isolated, in the sense of being free from contaminants such as antibodies able to bind other polypeptides and/or free of serum components. Monoclonal antibodies are preferred for some purposes, though polyclonal antibodies are within the scope of the present invention.

10

Antibodies may be obtained using techniques which are standard in the art. Methods of producing antibodies include immunising a mammal (e.g. mouse, rat, rabbit, horse, goat, sheep or monkey) with the protein or a fragment thereof. Antibodies may be obtained from immunised animals using any of a variety of techniques known in the art, and screened, preferably using binding of antibody to antigen of interest. For instance, Western blotting techniques or immunoprecipitation may be used (Armitage et al, Nature, 357:80-82, 1992). Isolation of antibodies and/or antibody-producing cells from an animal may be accompanied by a step of sacrificing the animal.

15

20

As an alternative or supplement to immunising a mammal with a peptide, an antibody specific for a protein may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which display functional immunoglobulin binding domains on their surfaces; for instance see WO92/01047. The library may be naive, that is constructed from sequences obtained from an organism which has not been immunised with any of the proteins (or fragments), or may be one constructed using sequences obtained from an organism which has been exposed to the antigen of interest.

25

30

Antibodies according to the present invention may be modified in a number of ways. Indeed the term "antibody" should be construed as covering any binding substance having a binding domain with the required specificity. Thus the invention covers antibody fragments, derivatives, functional equivalents and homologues of antibodies, including synthetic molecules and molecules whose shape mimics that of an antibody enabling it to bind an antigen or epitope.

35

40

Example antibody fragments, capable of binding an antigen or other binding partner are the Fab fragment consisting of the VL, VH, C1 and CH1 domains; the Fd fragment consisting of the VH and CH1 domains; the Fv fragment consisting of the VL and VH domains of a single arm of an

45

antibody; the dAb fragment which consists of a VH domain; isolated CDR regions and F(ab')<sub>2</sub> fragments, a bivalent fragment including two Fab fragments linked by a disulphide bridge at the hinge region. Single chain Fv fragments are also included.

5

Humanised antibodies in which CDRs from a non-human source are grafted onto human framework regions, typically with the alteration of some of the framework amino acid residues, to provide antibodies which are less immunogenic than the parent non-human antibodies, are also included within the present invention.

10

A hybridoma producing a monoclonal antibody according to the present invention may be subject to genetic mutation or other changes. It will further be understood by those skilled in the art that a monoclonal antibody can be subjected to the techniques of recombinant DNA technology to produce other antibodies or chimeric molecules which retain the specificity of the original antibody. Such techniques may involve introducing DNA encoding the immunoglobulin variable region, or the complementarity determining regions (CDRs), of an antibody to the constant regions, or constant regions plus framework regions, of a different immunoglobulin. See, for instance, EP-A-184187, GB-A-2188638 or EP-A-0239400. Cloning and expression of chimeric antibodies are described in EP-A-0120694 and EP-A-0125023.

15

20

25

Hybridomas capable of producing antibody with desired binding characteristics are within the scope of the present invention, as are host cells, eukaryotic or prokaryotic, containing nucleic acid encoding antibodies (including antibody fragments) and capable of their expression. The invention also provides methods of production of the antibodies including growing a cell capable of producing the antibody under conditions in which the antibody is produced, and preferably secreted.

30

35

The reactivities of antibodies on a sample may be determined by any appropriate means. Tagging with individual reporter molecules is one possibility. The reporter molecules may directly or indirectly generate detectable, and preferably measurable, signals. The linkage of reporter molecules may be directly or indirectly, covalently, e.g. via a peptide bond or non-covalently. Linkage via a peptide bond may be as a result of recombinant expression of a gene fusion encoding antibody and reporter molecule.

40

45

One favoured mode is by covalent linkage of each antibody with an individual fluorochrome, phosphor or laser dye with spectrally isolated absorption or emission characteristics. Suitable fluorochromes include

fluorescein, rhodamine, phycoerythrin and Texas Red. Suitable chromogenic dyes include diaminobenzidine.

5 Other reporters include macromolecular colloidal particles or particulate material such as latex beads that are coloured, magnetic or paramagnetic, and biologically or chemically active agents that can directly or indirectly cause detectable signals to be visually observed, electronically detected or otherwise recorded. These molecules may be enzymes which catalyse reactions that develop or  
10 change colours or cause changes in electrical properties, for example. They may be molecularly excitable, such that electronic transitions between energy states result in characteristic spectral absorptions or emissions. They may include chemical entities used in conjunction with biosensors. Biotin/avidin or biotin/streptavidin and alkaline  
15 phosphatase detection systems may be employed.

The mode of determining binding is not a feature of the present invention and those skilled in the art are able to choose a suitable mode according to their preference and general knowledge.

20 Antibodies according to the present invention may be used in screening for the presence of a polypeptide, for example in a test sample containing cells or cell lysate as discussed, and may be used in purifying and/or isolating a polypeptide according to the present invention, for instance following production of the polypeptide by  
25 expression from encoding nucleic acid therefor. Antibodies may modulate the activity of the polypeptide to which they bind and so, if that polypeptide has a deleterious effect in an individual, may be useful in a therapeutic context (which may include prophylaxis).

30 An antibody may be provided in a kit, which may include instructions for use of the antibody, e.g. in determining the presence of a particular substance in a test sample. One or more other reagents may be included, such as labelling molecules, buffer solutions, elutants and so on. Reagents may be provided within containers which protect  
35 them from the external environment, such as a sealed vial.

#### Diagnostic Methods

40 A number of methods are known in the art for analysing biological samples from individuals to determine whether the individual carries a BRCA2 allele predisposing them to cancer, especially breast cancer (female or male) or ovarian cancer. The purpose of such analysis may be used for diagnosis or prognosis, and serve to detect the presence of  
45 an existing cancer, to help identify the type of cancer, to assist a



physician in determining the severity or likely course of the cancer and/or to optimise treatment of it. Alternatively, the methods can be used to detect BRCA2 alleles that are statistically associated with a susceptibility to cancer in the future, e.g. early onset breast cancer, identifying individuals who would benefit from regular screening to provide early diagnosis of cancer. Examples of methods of screening for BRCA2 mutations are set out in example 5 below.

Broadly, the methods divide into those screening for the presence of BRCA2 nucleic acid sequences or alleles or variants thereof, and those that rely on detecting the presence or absence of the BRCA2 polypeptide. The methods make use of biological samples from individuals that are suspected of contain the nucleic acid sequences or polypeptide. Examples of biological samples include blood, plasma, serum, tissue samples, tumour samples, saliva and urine.

Exemplary approaches for detecting BRCA2 nucleic acid or polypeptides include:

(a) comparing the sequence of nucleic acid in the sample with the BRCA2 nucleic acid sequence to determine whether the sample from the patient contains mutations; or,

(b) determining the presence in a sample from a patient of the polypeptide encoded by the BRCA2 gene and, if present, determining whether the polypeptide is full length, and/or is mutated, and/or is expressed at the normal level; or,

(c) using DNA fingerprinting to compare the restriction pattern produced when a restriction enzyme cuts a sample of nucleic acid from the patient with the restriction pattern obtained from normal BRCA2 gene or from known mutations thereof; or,

(d) using a specific binding member capable of binding to a BRCA2 nucleic acid sequence (either a normal sequence or a known mutated sequence), the specific binding member comprising nucleic acid hybridisable with the BRCA2 sequence, or substances comprising an antibody domain with specificity for a native or mutated BRCA2 nucleic acid sequence or the polypeptide encoded by it, the specific binding member being labelled so that binding of the specific binding member to its binding partner is detectable; or,

(e) using PCR involving one or more primers based on normal or mutated BRCA2 gene sequence to screen for normal or mutant BRCA2 gene in a sample from a patient.

A "specific binding pair" comprises a specific binding member (sbm) and a binding partner (bp) which have a particular specificity for each other and which in normal conditions bind to each other in preference to other molecules. Examples of specific binding pairs are antigens

and antibodies, molecules and receptors and complementary nucleotide sequences. The skilled person will be able to think of many other examples and they do not need to be listed here. Further, the term "specific binding pair" is also applicable where either or both of the specific binding member and the binding partner comprise a part of a larger molecule. In embodiments in which the specific binding pair are nucleic acid sequences, they will be of a length to hybridise to each other under the conditions of the assay, preferably greater than 10 nucleotides long, more preferably greater than 15 or 20 nucleotides long.

In most embodiments for screening for BRCA2 susceptibility alleles, the BRCA2 nucleic acid in the sample will initially be amplified, e.g. using PCR, to increase the amount of the analyte as compared to other sequences present in the sample. This allows the target sequences to be detected with a high degree of sensitivity if they are present in the sample. This initial step may be avoided by using highly sensitive array techniques that are becoming increasingly important in the art.

The identification of the BRCA2 gene and its association with cancer paves the way for aspects of the present invention to provide the use of materials and methods, such as are disclosed and discussed above, for establishing the presence or absence in a test sample of an variant form of the gene, in particular an allele or variant specifically associated with cancer, especially breast and ovarian cancer. This may be for diagnosing a predisposition of an individual to cancer. It may be for diagnosing cancer of a patient with the disease as being associated with the gene.

This allows for planning of appropriate therapeutic and/or prophylactic treatment, permitting stream-lining of treatment by targeting those most likely to benefit.

A variant form of the gene may contain one or more insertions, deletions, substitutions and/or additions of one or more nucleotides compared with the wild-type sequence (such as shown in table 1) which may or may not disrupt the gene function. Differences at the nucleic acid level are not necessarily reflected by a difference in the amino acid sequence of the encoded polypeptide. However, a mutation or other difference in a gene may result in a frame-shift or stop codon, which could seriously affect the nature of the polypeptide produced (if any), or a point mutation or gross mutational change to the encoded polypeptide, including insertion, deletion, substitution and/or addition of one or more amino acids or regions in the polypeptide. A mutation in a promoter sequence or other regulatory region may prevent

or reduce expression from the gene or affect the processing or stability of the mRNA transcript.

5 There are various methods for determining the presence or absence in a test sample of a particular nucleic acid sequence, such as the sequence shown in figures 1, 2, 4 or 7 or a variant or allele thereof.

10 Tests may be carried out on preparations containing genomic DNA, cDNA and/or mRNA. Testing cDNA or mRNA has the advantage of the complexity of the nucleic acid being reduced by the absence of intron sequences, but the possible disadvantage of extra time and effort being required in making the preparations. RNA is more difficult to manipulate than DNA because of the wide-spread occurrence of RN'ases.

15 Nucleic acid in a test sample may be sequenced and the sequence compared with the sequence shown in figures 1, 2, 4 or 7, to determine whether or not a difference is present. If so, the difference can be compared with known susceptibility alleles (e.g. as summarised in table 1) to determine whether the test nucleic acid contains one or more of  
20 the variations indicated, or the difference can be investigated for association with cancer.

Since it will not generally be time- or labour-efficient to sequence all nucleic acid in a test sample or even the whole BRCA2 gene, a  
25 specific amplification reaction such as PCR using one or more pairs of primers may be employed to amplify the region of interest in the nucleic acid, for instance the BRCA2 gene or a particular region in which mutations associated with cancer susceptibility occur. Exemplary primers for this purpose are shown in figure 8. The amplified nucleic  
30 acid may then be sequenced as above, and/or tested in any other way to determine the presence or absence of a particular feature. Nucleic acid for testing may be prepared from nucleic acid removed from cells or in a library using a variety of other techniques such as restriction enzyme digest and electrophoresis.

35 Nucleic acid may be screened using a variant- or allele-specific probe. Such a probe corresponds in sequence to a region of the BRCA2 gene, or its complement, containing a sequence alteration known to be associated with cancer susceptibility. Under suitably stringent conditions,  
40 specific hybridisation of such a probe to test nucleic acid is indicative of the presence of the sequence alteration in the test nucleic acid. For efficient screening purposes, more than one probe may be used on the same test sample.

45 Allele- or variant-specific oligonucleotides may similarly be used in

5 PCR to specifically amplify particular sequences if present in a test sample. Assessment of whether a PCR band contains a gene variant may be carried out in a number of ways familiar to those skilled in the art. The PCR product may for instance be treated in a way that enables one to display the mutation or polymorphism on a denaturing polyacrylamide DNA sequencing gel, with specific bands that are linked to the gene variants being selected.

10 An alternative or supplement to looking for the presence of variant sequences in a test sample is to look for the presence of the normal sequence, e.g. using a suitably specific oligonucleotide probe or primer.

15 Use of oligonucleotide probes and primers has been discussed in more detail above.

20 Approaches which rely on hybridisation between a probe and test nucleic acid and subsequent detection of a mismatch may be employed. Under appropriate conditions (temperature, pH etc.), an oligonucleotide probe will hybridise with a sequence which is not entirely complementary. The degree of base-pairing between the two molecules will be sufficient for them to anneal despite a mis-match. Various approaches are well known in the art for detecting the presence of a mis-match between two annealing nucleic acid molecules.

25 For instance, RN'ase A cleaves at the site of a mis-match. Cleavage can be detected by electrophoresing test nucleic acid to which the relevant probe or probe has annealed and looking for smaller molecules (i.e. molecules with higher electrophoretic mobility) than the full length probe/test hybrid. Other approaches rely on the use of enzymes such as resolvases or endonucleases.

30 Thus, an oligonucleotide probe that has the sequence of a region of the normal BRCA2 gene (either sense or anti-sense strand) in which mutations associated with cancer susceptibility are known to occur (e.g. see table 1) may be annealed to test nucleic acid and the presence or absence of a mis-match determined. Detection of the presence of a mis-match may indicate the presence in the test nucleic acid of a mutation associated with cancer susceptibility. On the other hand, an oligonucleotide probe that has the sequence of a region of the BRCA2 gene including a mutation associated with cancer susceptibility may be annealed to test nucleic acid and the presence or absence of a mis-match determined. The absence of a mis-match may indicate that the nucleic acid in the test sample has the normal sequence. In either case, a battery of probes to different regions of the gene may be

employed.

5 The presence of differences in sequence of nucleic acid molecules may be detected by means of restriction enzyme digestion, such as in a method of DNA fingerprinting where the restriction pattern produced when one or more restriction enzymes are used to cut a sample of nucleic acid is compared with the pattern obtained when a sample containing the normal gene or a variant or allele is digested with the same enzyme or enzymes.

10 The presence of absence of a lesion in a promoter or other regulatory sequence may also be assessed by determining the level of mRNA production by transcription or the level of polypeptide production by translation from the mRNA.

15 A test sample of nucleic acid may be provided for example by extracting nucleic acid from cells, e.g. in saliva or preferably blood, or for pre-natal testing from the amnion, placenta or foetus itself.

20 There are various methods for determining the presence or absence in a test sample of a particular polypeptide, such as the polypeptide with the amino acid sequence shown in figure 3, 5 or 7 or an amino acid sequence variant or allele thereof.

25 A sample may be tested for the presence of a binding partner for a specific binding member such as an antibody (or mixture of antibodies), specific for one or more particular variants of the polypeptide shown in figures 3, 5 or 7.

30 A sample may be tested for the presence of a binding partner for a specific binding member such as an antibody (or mixture of antibodies), specific for the polypeptide shown in figures 3, 5 or 7.

35 In such cases, the sample may be tested by being contacted with a specific binding member such as an antibody under appropriate conditions for specific binding, before binding is determined, for instance using a reporter system as discussed. Where a panel of antibodies is used, different reporting labels may be employed for each antibody so that binding of each can be determined.

40 A specific binding member such as an antibody may be used to isolate and/or purify its binding partner polypeptide from a test sample, to allow for sequence and/or biochemical analysis of the polypeptide to determine whether it has the sequence and/or properties of the polypeptide whose sequence is shown in figures 3, 5 or 7, or if it is  
45

a mutant or variant form. Amino acid sequence is routine in the art using automated sequencing machines.

5 There is also an increasing tendency in the diagnostic field towards miniaturisation of such assays, e.g. making use of binding agents (such as antibodies or nucleic acid sequences) immobilised in small, discrete locations (microspots) and/or as arrays on solid supports or on diagnostic chips. These approaches can be particularly valuable as they can provide great sensitivity (particularly through the use of  
10 fluorescently labelled reagents), require only very small amounts of biological sample from individuals being tested and allow a variety of separate assays can be carried out simultaneously. This latter advantage can be useful as it provides an assay for different mutations in the BRCA2 gene or another cancer susceptibility gene (such as BRCA1, see EP-A-705902) to be carried out using a single sample. Examples of techniques enabling this miniaturised technology are provided in WO84/01031, WO88/1058, WO89/01157, WO93/8472, WO95/18376/ WO95/18377, WO95/24649 and EP-A-0373203. Thus, in a further aspect, the present  
20 invention provides a kit comprising a support or diagnostic chip having immobilised thereon one or more binding agents capable of specifically binding BRCA2 nucleic acid or polypeptides, optionally in combination with other reagents (such as labelled developing reagents) needed to carrying out an assay.

## 25 Therapeutics

### Pharmaceuticals and Peptide Therapies

30 The BRCA2 polypeptides, antibodies, peptides and nucleic acid of the invention can be formulated in pharmaceutical compositions. These compositions may comprise, in addition to one of the above substances, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the  
35 active ingredient. The precise nature of the carrier or other material may depend on the route of administration, e.g. oral, intravenous, cutaneous or subcutaneous, nasal, intramuscular, intraperitoneal routes.

40 Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant. Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline  
45 solution, dextrose or other saccharide solution or glycols such as

ethylene glycol, propylene glycol or polyethylene glycol may be included.

5 For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as sodium chloride injection, Ringer's  
10 injection, lactated Ringer's injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

15 Whether it is a polypeptide, antibody, peptide, nucleic acid molecule, small molecule or other pharmaceutically useful compound according to the present invention that is to be given to an individual, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show  
20 benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors, and typically takes account of  
25 the disorder to be treated, the condition of the individual patient, the site of delivery, the method of administration and other factors known to practitioners. Examples of the techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.

30 Alternatively, targeting therapies may be used to deliver the active agent more specifically to certain types of cell, by the use of targeting systems such as antibody or cell specific ligands. Targeting may be desirable for a variety of reasons; for example if the agent is  
35 unacceptably toxic, or if it would otherwise require too high a dosage, or if it would not otherwise be able to enter the target cells.

40 Instead of administering these agents directly, they could be produced in the target cells by expression from an encoding gene introduced into the cells, eg in a viral vector (a variant of the VDEPT technique - see below). The vector could be targeted to the specific cells to be treated, or it could contain regulatory elements which are switched on more or less selectively by the target cells.

45 Alternatively, the agent could be administered in a precursor form, for

conversion to the active form by an activating agent produced in, or targeted to, the cells to be treated. This type of approach is sometimes known as ADEPT or VDEPT; the former involving targeting the activating agent to the cells by conjugation to a cell-specific antibody, while the latter involves producing the activating agent, eg an enzyme, in a vector by expression from encoding DNA in a viral vector (see for example, EP-A-415731 and WO90/07936).

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

#### Methods of Gene Therapy

As a further alternative, the nucleic acid encoded the authentic biologically active BRCA2 polypeptide could be used in a method of gene therapy, to treat a patient who is unable to synthesize the active polypeptide or unable to synthesize it at the normal level, thereby providing the effect provided by wild-type BRCA2 and suppressing the occurrence of cancer and/or reduce the size or extent of existing cancer in the target cells.

Vectors such as viral vectors have been used in the prior art to introduce genes into a wide variety of different target cells. Typically the vectors are exposed to the target cells so that transfection can take place in a sufficient proportion of the cells to provide a useful therapeutic or prophylactic effect from the expression of the desired polypeptide. The transfected nucleic acid may be permanently incorporated into the genome of each of the targeted tumour cells, providing long lasting effect, or alternatively the treatment may have to be repeated periodically.

A variety of vectors, both viral vectors and plasmid vectors, are known in the art, see US Patent No. 5,252,479 and WO93/07282. In particular, a number of viruses have been used as gene transfer vectors, including papovaviruses, such as SV40, vaccinia virus, herpesviruses, including HSV and EBV, and retroviruses. Many gene therapy protocols in the prior art have used disabled murine retroviruses.

As an alternative to the use of viral vectors other known methods of introducing nucleic acid into cells includes electroporation, calcium phosphate co-precipitation, mechanical techniques such as microinjection, transfer mediated by liposomes and direct DNA uptake and receptor-mediated DNA transfer.



As mentioned above, the aim of gene therapy using nucleic acid encoding the BRCA2 polypeptide, or an active portion thereof, is to increase the amount of the expression product of the nucleic acid in cells in which the level of the wild-type BRCA2 polypeptide is absent or present only at reduced levels. Such treatment may be therapeutic in the treatment of cells which are already cancerous or prophylactic in the treatment of individuals known through screening to have a BRCA2 susceptibility allele and hence a predisposition to cancer.

Gene transfer techniques which selectively target the BRCA2 nucleic acid to breast and/or ovarian tissues are preferred. Examples of this included receptor-mediated gene transfer, in which the nucleic acid is linked to a protein ligand via polylysine, with the ligand being specific for a receptor present on the surface of the target cells.

Antisense technology based on the BRCA2 nucleic acid sequences is discussed above.

#### Methods of Screening for Drugs

A polypeptide according to the present invention may be used in screening for molecules which affect or modulate its activity or function. Such molecules may be useful in a therapeutic (possibly including prophylactic) context.

It is well known that pharmaceutical research leading to the identification of a new drug may involve the screening of very large numbers of candidate substances, both before and even after a lead compound has been found. This is one factor which makes pharmaceutical research very expensive and time-consuming. Means for assisting in the screening process can have considerable commercial importance and utility. Such means for screening for substances potentially useful in treating or preventing cancer is provided by polypeptides according to the present invention. Substances identified as modulators of the polypeptide represent an advance in the fight against cancer since they provide basis for design and investigation of therapeutics for in vivo use.

A method of screening for a substance which modulates activity of a polypeptide may include contacting one or more test substances with the polypeptide in a suitable reaction medium, testing the activity of the treated polypeptide and comparing that activity with the activity of the polypeptide in comparable reaction medium untreated with the test substance or substances. A difference in activity between the treated and untreated polypeptides is indicative of a modulating effect of the

relevant test substance or substances.

Combinatorial library technology provides an efficient way of testing a potentially vast number of different substances for ability to modulate activity of a polypeptide. Such libraries and their use are known in the art. The use of peptide libraries is preferred.

Prior to or as well as being screened for modulation of activity, test substances may be screened for ability to interact with the polypeptide, e.g. in a yeast two-hybrid system (which requires that both the polypeptide and the test substance can be expressed in yeast from encoding nucleic acid). This may be used as a coarse screen prior to testing a substance for actual ability to modulate activity of the polypeptide. Alternatively, the screen could be used to screen test substances for binding to a BRCA2 specific binding partner, to find mimetics of the BRCA2 polypeptide, e.g. for testing as cancer therapeutics.

Following identification of a substance which modulates or affects polypeptide activity, the substance may be investigated further. Furthermore, it may be manufactured and/or used in preparation, i.e. manufacture or formulation, of a composition such as a medicament, pharmaceutical composition or drug. These may be administered to individuals.

Thus, the present invention extends in various aspects not only to a substance identified using a nucleic acid molecule as a modulator of polypeptide activity, in accordance with what is disclosed herein, but also a pharmaceutical composition, medicament, drug or other composition comprising such a substance, a method comprising administration of such a composition to a patient, e.g. for treatment (which may include preventative treatment) of cancer, use of such a substance in manufacture of a composition for administration, e.g. for treatment of cancer, and a method of making a pharmaceutical composition comprising admixing such a substance with a pharmaceutically acceptable excipient, vehicle or carrier, and optionally other ingredients.

A substance identified using as a modulator of polypeptide function may be peptide or non-peptide in nature. Non-peptide "small molecules" are often preferred for many in vivo pharmaceutical uses. Accordingly, a mimetic or mimic of the substance (particularly if a peptide) may be designed for pharmaceutical use.

The designing of mimetics to a known pharmaceutically active compound

is a known approach to the development of pharmaceuticals based on a "lead" compound. This might be desirable where the active compound is difficult or expensive to synthesise or where it is unsuitable for a particular method of administration, e.g. peptides are unsuitable active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary canal. Mimetic design, synthesis and testing is generally used to avoid randomly screening large number of molecules for a target property.

There are several steps commonly taken in the design of a mimetic from a compound having a given target property. Firstly, the particular parts of the compound that are critical and/or important in determining the target property are determined. In the case of a peptide, this can be done by systematically varying the amino acid residues in the peptide, eg by substituting each residue in turn. Alanine scans of peptide are commonly used to refine such peptide motifs. These parts or residues constituting the active region of the compound are known as its "pharmacophore".

Once the pharmacophore has been found, its structure is modelled to according its physical properties, eg stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g. spectroscopic techniques, X-ray diffraction data and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of a pharmacophore, rather than the bonding between atoms) and other techniques can be used in this modelling process.

In a variant of this approach, the three-dimensional structure of the ligand and its binding partner are modelled. This can be especially useful where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the design of the mimetic.

A template molecule is then selected onto which chemical groups which mimic the pharmacophore can be grafted. The template molecule and the chemical groups grafted on to it can conveniently be selected so that the mimetic is easy to synthesise, is likely to be pharmacologically acceptable, and does not degrade in vivo, while retaining the biological activity of the lead compound. Alternatively, where the mimetic is peptide based, further stability can be achieved by cyclising the peptide, increasing its rigidity. The mimetic or mimetics found by this approach can then be screened to see whether they have the target property, or to what extent they exhibit it. Further optimisation or modification can then be carried out to arrive at one or more final mimetics for in vivo or clinical testing.

Screening for Substances Affecting BRCA2 Expression

5 The present invention also provides the use of all or part of the nucleic acid sequence of the BRCA2 promoter region shown in figure 6 in methods of screening for substances which modulate the activity of the promoter and increase or decrease the level of BRCA2 expression.

10 "Promoter activity" is used to refer to ability to initiate transcription. The level of promoter activity is quantifiable for instance by assessment of the amount of mRNA produced by transcription from the promoter or by assessment of the amount of protein product produced by translation of mRNA produced by transcription from the promoter. The amount of a specific mRNA present in an expression system may be determined for example using specific oligonucleotides  
15 which are able to hybridise with the mRNA and which are labelled or may be used in a specific amplification reaction such as the polymerase chain reaction. Use of a reporter gene facilitates determination of promoter activity by reference to protein production.

20 Further provided by the present invention is a nucleic acid construct comprising a BRCA2 promoter region set out in figure 6 or a fragment, mutant, allele, derivative or variant thereof able to promoter transcription, operably linked to a heterologous gene, e.g. a coding sequence. A "heterologous" or "exogenous" gene is generally not a modified form of BRCA2. Generally, the gene may be transcribed into mRNA which may be translated into a peptide or polypeptide product  
25 which may be detected and preferably quantitated following expression. A gene whose encoded product may be assayed following expression is termed a "reporter gene", i.e. a gene which "reports" on promoter activity.  
30

The reporter gene preferably encodes an enzyme which catalyses a reaction which produces a detectable signal, preferably a visually detectable signal, such as a coloured product. Many examples are  
35 known, including  $\beta$ -galactosidase and luciferase.  $\beta$ -galactosidase activity may be assayed by production of blue colour on substrate, the assay being by eye or by use of a spectrophotometer to measure absorbance. Fluorescence, for example that produced as a result of luciferase activity, may be quantitated using a spectrophotometer.  
40 Radioactive assays may be used, for instance using chloramphenicol acetyltransferase, which may also be used in non-radioactive assays. The presence and/or amount of gene product resulting from expression from the reporter gene may be determined using a molecule able to bind the product, such as an antibody or fragment thereof. The binding  
45 molecule may be labelled directly or indirectly using any standard

technique.

Those skilled in the art are well aware of a multitude of possible reporter genes and assay techniques which may be used to determine gene activity. Any suitable reporter/assay may be used and it should be appreciated that no particular choice is essential to or a limitation of the present invention.

Nucleic acid constructs comprising a promoter (as disclosed herein) and a heterologous gene (reporter) may be employed in screening for a substance able to modulate activity of the promoter. For therapeutic purposes, e.g. for treatment of cancer a substance able to up-regulate expression of the promoter directing the expression of normal may be sought. Alternatively, substances to down-regulate the promoter may help to prevent or inhibit the production of mutated BRCA2 polypeptide, if this is an agent implicated in the development of cancer. A method of screening for ability of a substance to modulate activity of a promoter may comprise contacting an expression system, such as a host cell, containing a nucleic acid construct as herein disclosed with a test or candidate substance and determining expression of the heterologous gene.

The level of expression in the presence of the test substance may be compared with the level of expression in the absence of the test substance. A difference in expression in the presence of the test substance indicates ability of the substance to modulate gene expression. An increase in expression of the heterologous gene compared with expression of another gene not linked to a promoter as disclosed herein indicates specificity of the substance for modulation of the promoter.

A promoter construct may be introduced into a cell line using any technique previously described to produce a stable cell line containing the reporter construct integrated into the genome. The cells may be grown and incubated with test compounds for varying times. The cells may be grown in 96 well plates to facilitate the analysis of large numbers of compounds. The cells may then be washed and the reporter gene expression analysed. For some reporters, such as luciferase the cells will be lysed then analysed.

Following identification of a substance which modulates or affects promoter activity, the substance may be investigated further. Furthermore, it may be manufactured and/or used in preparation, i.e. manufacture or formulation, of a composition such as a medicament, pharmaceutical composition or drug. These may be administered to

individuals.

Construction of Animal Models for BRCA2 Deficiency

5 The construction of animal models for BRCA2 deficiency can be carried out using standard techniques for introducing mutations into, for example, a mouse germ-line. In one example of this approach, a vector carrying an insertional mutation within exon 11 of the mouse BRCA2 gene is transfected into embryonic stem cells. Clones in which the mutant  
10 version of the gene has replaced with wild type are identified by Southern blot hybridisation. The clones are then amplified and cells are injected into mouse blastocyst stage embryos. Mice in which the injected cells have contributed to the development of the mouse are identified by Southern blotting. These chimeric mice are then bred to  
15 produce mice which carry one copy of the mutation in the germ line. These heterozygous mutant animals are then bred to produce mice carrying mutations in the BRCA2 gene homozygously. The mice having a heterozygous mutation in BRCA2 may be a suitable model for human individuals having one copy of the gene mutated in the germ line who  
20 are at high risk of developing breast cancer.

Example 1 Identification of the BRCA2 Gene

25 Following the definition of the interval D13S289-D13S267 in which the BRCA2 gene is believed to be located (Wooster et al (1994)), the following set of procedures were used identify the gene itself. A yeast artificial chromosome (YAC) contig was constructed of YACs believed to be in the region from the CEPH database. Chimerism of YACs was examined using fluorescent in situ hybridisation. Overlaps between  
30 YACs were established by amplification of sequence tagged sites (STS), hybridisation of STS PCR products and alu-PCR fingerprinting.

Using probes localised to and derived from the YAC contig, a P1 artificial chromosome (PAC) library was screened and positive PACs  
35 isolated. PACs were rehybridised to the PAC library to fill in gaps between clones. A PAC contig over a 1 megabase region was then assembled.

Overlaps between PACs were identified by:

- 40 (a) amplification of STS;  
(b) hybridisation of end probes produced by linear PCR;  
(c) HindIII/Sau3A fingerprinting.

45 Additional polymorphic markers from the region were identified by screening M13 libraries constructed from PACs, with oligonucleotides

containing repetitive sequences that are commonly polymorphic (GTn, GAn, CAGn, GATAn, GAATn). Using new markers, the region in which BRCA2 is likely to be located was narrowed further.

- 5 Transcripts from the region were identified by use of PAC DNAs in:
- (a) exon trapping experiments using a lambda GET vector;
  - (b) hybrid selection between PAC genomic DNA and cDNA libraries.

Exons and cDNA fragments identified by these methods were sequenced.

10

Using primers synthesised from these sequences as probes, 15kb genomic subclones of the PACs that carry all or part of the cDNAs and exons were identified. These were sequenced from the cDNA sequence out into adjacent flanking genomic sequences.

15

Using oligonucleotide primers synthesised on the basis of these flanking genomic sequences, genomic DNA fragments containing the potential transcript were amplified and screened for mutations through the following DNA samples; DNAs from 46 families that show evidence of linkage to BRCA2 and/or show evidence against linkage to BRCA1 and/or do not have BRCA1 mutations and/or do have evidence of male breast cancer. Also screened were cancer cell lines that are homozygous for all polymorphic markers through the BRCA2 region; DNAs from 12 primary human tumour DNAs that show evidence of loss of heterozygosity in the BRCA2 region.

25

Sequence variants were detected by running <sup>32</sup>P labelled amplification products through:

- 30 (a) non denaturing polyacrylamide gels, at room temperature and at 4°C;
- (b) denaturing polyacrylamide gels.

35

Fragments that showed altered mobility were reamplified using the PCR and directly sequenced. Sequences were run on an ABI 377 DNA sequencer.

40

In the course of this work, several sequence variants were detected, most of which are believed to be non disease associated polymorphisms. However, an abnormality detected in one fragment was predicted to destroy a splice site and create a termination codon in a family that shows strong linkage to BRCA2. The variant segregates with the disease chromosome. This is precisely the type of abnormality that would be expected within a cancer susceptibility gene and made this fragment a strong candidate for part of the BRCA2 gene.

45

This fragment was then used (a) as a probe in screening of cDNA libraries, (b) starting sequence for PCR amplification experiments from cDNAs and cDNA libraries. Positive clones and amplification fragments were sequenced. These results are shown in the figures 1 and 2.

5 From the initial isolation of this part of exon 16 of the BRCA2 gene, a variety of conventional techniques were then used to isolate the remaining portions of the BRCA2 sequence. The 900,000bp sequence released on the Internet (the Sanger sequence) assisted in this  
10 procedure. Figure 4 shows around 75% of the cDNA sequence isolated by the inventors, while the complete BRCA2 gene sequence, including the exon/intron structure, is shown in figure 7. The promoter region of the gene is set out in figure 6. These sequences could be downloaded from the ftp web sites to a local computer. The sequences can then be  
15 analysed using programs such as BLAST, FASTA, GRAIL or GeneFinder to identify putative coding regions. Oligos can then be designed to these predicted coding regions and these would be used in RT-PCR to confirm or refute the coding regions. The coding regions would also be compared with the experimental results obtained from the procedures set  
20 out herein.

Thus, given knowledge of a part of the BRCA2 sequence and its orientation within the Sanger sequence, the skilled person could readily isolate the rest of the sequence using a conventional exon  
25 prediction program to localise potential exons/open reading frames in the same orientation as the known portion of the BRCA2 sequence, and then do exon connection in which potential exons are used to design primers that sit in putative coding sequence. The primers could be designed in such a way as to have a primer based on known BRCA2  
30 sequence and an unknown primer, such that when these are used to prime from cDNA, a product contiguous with known sequence would be produced if the unknown primer is a part of the BRCA2 gene. This process could be continued in an iterative manner, readily allowing the skilled person to walk through the Sanger sequence to obtain the full BRCA2  
35 sequence, obtaining the intron/exon boundaries as part of the procedure. In the case of BRCA2, a large portion of the coding sequence (about 5kb) is in exon 11, part of which is included in the 10% of the sequence disclosed in figures 1 and 2.

40 By way of reference, the techniques described above and others useful in the isolation of genes by positional cloning are reviewed in Monaco, Curr. Opin. Gen. Devel., 4:360-365, 1994 and summarised in EP-A-0705902.

45 By way of further assistance, a protocol for the isolation of the full



length sequence and the polypeptide encoded by the BRCA2 gene from the sequence shown in figures 1, 2 and 4 is set out below, and could be used iteratively to isolate successive portions of the BRCA2 gene.

5      **Screening cDNA libraries by hybridisation.**

Presently identified cDNA fragments can be <sup>32</sup>P labelled and hybridised to various widely available plated or gridded cDNA libraries. Positive clones can then be isolated, and subject to replating and rehybridisation if necessary until a pure clone has been isolated. DNA  
10      can then be made from pure clones and will be sequenced by conventional Sanger dideoxy sequencing on a ABI 377 DNA sequencer.

**Screening cDNA libraries by PCR amplification.**

Oligonucleotides based on sequences within the BRCA2 sequences  
15      disclosed herein can be used in conjunction with oligonucleotides designed to prime from the cloning vector in PCR amplifications of aliquots of widely available cDNA libraries. This will allow amplification of fragments of the BRCA2 cDNA positioned between the currently known fragment and the cloning insertion site. Products of  
20      the PCR amplification can then be sequenced using Sanger dideoxy sequencing on an ABI 377 sequencer.

**Rapid amplification of cDNA ends (RACE).**

Primary cDNAs synthesised from a number of different tissue RNAs can be  
25      ligated to an oligonucleotide linker. After purification, PCR amplifications can be performed using an oligonucleotide that primes from the current BRCA2 cDNA sequence and a second oligonucleotide that primes from the linker. Amplification products will be directly sequenced using Sanger dideoxy sequencing. RACE is described in (6).  
30

The new sequences can then be integrated into the full sequence of the gene by detection of overlaps with previously known components of the sequence.

35      The screening of cDNA or genomic libraries with selected probes can be conducted using standard procedures, for instance as described in Ausubel et al and Sambrook et al (supra).

40      These techniques allow the full coding sequence to be isolated based on the information disclosed in figures 1, 2 and 4. The full length sequence is defined as the sequence between a translation initiation codon (ATG) and a translation termination codon (TAA, TAG, TGA) between which there is an open reading frame. This in turn can be used to define the intron-exon structure of the gene. Primers can then be  
45      designed to flank each exon so that the whole coding sequence of the

gene can be amplified from genomic DNA, see for example the primers disclosed in figure 8 or in Nature Genetics, 12:333-337, 1996.

5 Further fragments of coding sequence were then amplified from genomic DNA in the previously described mutation testing set in order to detect additional disease associated mutations in the BRCA2 gene.

Example 2 Mutations in the BRCA2 Gene  
Identification of Six Mutations in the BRCA2 Gene

10 In a first study, the inventors found a series of mutations in the BRCA2 sequence by comparing the native sequence with sequences obtained from families with a history of multiple cases of early onset breast cancer. The locations of these mutations in the amino acid sequence  
15 are shown by boxing the residues in the native sequence which are affected (see figure 5). The mutations to the BRCA2 gene are summarised in table 3, which shows the families in the left hand column against mutations in the BRCA2 gene in the right hand column. The remaining columns in table 3 from left to right are as follows:

20 Number of FBCs = Number of female breast cancers in the family.

25 Number of FBCs<50 = Number of female breast cancers occurring in individuals under 50 years of age.

Number of OvCs = Number of ovarian cancers in the family.

30 Number of MBCs = Number of male breast cancers.

35 The columns referring to LOD scores at BRCA1 and BRCA2 show the chance that the incidence of the cancers in a family is linked to the BRCA1 or BRCA2 genes, with a more positive value indicating a greater chance of linkage. Thus, for example, a LOD score of +3 indicates a strong linkage between the incidence of cancer and the given gene in that family.

40 To identify BRCA2, genomic DNA fragments of less than 300bp containing putative coding sequences were screened for mutations. At least one affected member of 46 breast cancer families was examined. Each family included in this set either shows evidence of linkage to BRCA2, and/or shows evidence of breast cancer. The majority, but probably not all, of these families would be expected to be due to BRCA2 mutations.

45

Disease associated mutations in most known cancer susceptibility genes usually result in truncation of the encoded protein and inactivation of critical functions. In the course of the mutational screen of candidate coding sequences from the BRCA2 region, the first detected sequence variant that was predicted to disrupt translation of an encoded protein was observed in IARC 2932. This family is clearly linked to BRCA2 with a multipoint LOD score of 3.01 using D13S260 and D13S267. A six base pair deletion removes the last five bases of the exon examined (exon S66), deletes the conserved G of the 5' splice site of the intron, and directly converts the codon TTT for phenylalanine to the termination codon TAA. By sequencing this mutation has been detected in lymphocyte DNA from two other early onset breast cancer cases in this family. The individuals examined share only the disease-associated haplotype. The mutation is absent in more than 500 chromosomes from normal individuals or in the remaining families and cancers. This finding therefore identified the candidate gene which was proved by the work described herein to be the BRCA2 gene.

To characterise the BRCA2 gene further, exon S66 was used to isolate a series of cDNA clones which represented segments of the BRCA2 candidate. From alignment of the cDNA and genomic sequence data, the candidate BRCA2 gene was found to lie in three sequence contigs which also contained other previously isolated transcribed sequences. The exon and open reading frame prediction program Genemark was used to define putative additional 5' exons of the gene. Contiguity of the transcription unit was confirmed by RT-PCR on cDNA and sequence analysis. The availability of extensive sequence information at the cDNA and genomic level allowed mutational analysis of further coding regions of the putative BRCA2 gene in samples from breast cancer families.

A TG deletion (6819delTG) and a TT (6503delTT) deletion were detected in families CRC B196 and CRC B211 respectively (tables 1 and 2). In both families the mutation has been detected by sequencing other individuals with early onset breast cancer who share only the haplotype of 13q microsatellite markers that segregates with the disease.

Therefore, the mutations are on the disease associated chromosomes. A CT deletion was detected in family IARC 3594. This mutation has arisen within a short repetitive sequence (CTCTCT), a feature that is characteristic of deletion/insertion mutations in many genes and which is presumed to be due to slippage during DNA synthesis. Finally, a T deletion (6174delT) and an AAAC deletion have been found in Montreal 681 and 440 respectively. Both these families include a male breast cancer case and previous analyses have indicated that the large

majority of such families will have BRCA2 mutations. All these mutations are predicted to generate frame shifts leading to premature termination codons. None of the mutations have been found in chromosomes from over 500 healthy women and are therefore unlikely to be polymorphisms. The identification of several different germline mutations that truncate the encoded protein in breast cancer families that are highly likely to be due to BRCA2 strongly suggests that we have identified the BRCA2 gene. In particular, the 6174delT mutation from Ashkenazi Jewish family data is reproduced in other families examined and so may be useful in screening individuals for a susceptibility to cancer, especially male or female breast cancer or ovarian cancer.

Northern analysis has demonstrated that BRCA2 is encoded by a transcript of 10-12kb which is present in normal breast epithelial cells, placenta, and the breast cancer cell line MCF7. This suggests that our present contig of cDNAs covering approximately 9kb (including 1.6kb of 3' untranslated sequence) may not include the whole BRCA2 coding sequence. The known sequence of 2329 amino acids encoded by the BRCA2 gene does not show strong homology to sequences in the publicly available DNA or protein databases. The homology and motifs of BRCA2 found in an analysis of the corresponding genes in other species is described in example 6. However, some weak matches were detected including, intriguingly, a very weak similarity to the BRCA1 protein over a restricted region (amino acids 1394-1474 in BRCA1 and 1783-1863 in the portion of BRCA2 shown in figure 5).

#### Identification of Further Mutations in the BRCA2 Gene

##### **Patient material**

Families were identified which contained a minimum of either three cases of breast cancer and for which DNA samples were available from a least one affected individual. Complete or partial analysis of the BRCA1 gene identified those families with disease associated germline mutations. The remaining families were analysed throughout the complete BRCA2 coding sequence for germline mutations, which are shown in table 1 against the incidences of breast cancer (including early onset breast cancer) and ovarian cancer. Borderline ovarian cancer was not included. Confirmation of diagnosis was available from pathology reports or death certificates.

##### **Mutation analysis of BRCA2 in ovarian cancer families**

Seventy seven families with two or more first degree relatives with epithelial ovarian cancer were analysed for BRCA2 mutations. Mutation screening was performed using a combination of the protein truncation

test (PTT) and non-radioactive single-strand conformation analysis/heteroduplex analysis (SSCA/HA). PTT was performed from genomic DNA for exon 11 in all cases and for the entire coding region in those individuals in which RNA derived from lymphoblastoid cell lines was available to perform reverse-transcriptase PCR. Primers were designed to PCR amplify exon 11 or the complete coding sequence in overlapping fragments ranging in size from 10 to 1.3kb.

PTT was performed using the TNT rabbit reticulocyte lysate system (Promega) incorporating <sup>35</sup>S methionine (Amersham) for protein detection. Protein products were electrophoresed on 12-15% SDS polyacrylamide gels at 30-60mA for 12-16 hours. SDS-PAGE gels were fixed in 30% methanol/10% glacial acetic acid, dried and exposed to Kodak X-Omat film for 16-72 hours. The approximate location of the sequence alteration resulting in truncated protein variants was localised and the region sequenced to confirm the precise nucleotide alteration. SSCA/HA was performed on genomic DNA for coding exons 2-10 and 12-27. The 5' and 3' splice boundaries for exon 11 were also analysed. SSCA and HA variant conformers were sequenced as previously described (3) in order to characterise the precise nucleotide change. The ability to detect mutations identified by PTT using SSCA/HA was detectable using SSCA/HA on fragments ranging in size from 300-600bp. All primer sequences designed for PTT and SSCA/HA are available on request (e-mail sg200@cam.ac.uk).

#### Mutation analysis of BRCA2 in breast cancer families

##### Statistical methods

We tested for an association between mutation location and disease phenotype using a permutation argument similar to that used by Gayther et al (12). Test statistics were based on standard chi-squared statistics for the difference in the rate of ovarian cancers, as a proportion of all ovarian and breast cancers, confirmed by position. In the main analysis, the statistic X was calculated assuming an alternative hypothesis in which different rates applied for mutations in three different regions, i.e. before codon n1-n2 and after n2 (i.e. a 2 degree of freedom chi-squared). The three putative missense mutations were ignored in this analysis as their significance is unclear. The statistic X was maximised over all possible values of the cutpoints n1 and n2 giving the statistic Xm. The significance of this maximised chi-squared was then computed empirically by calculating the value of Xm in 50,000 datasets in which the 26 mutations were randomly permuted among the families. We also computed a statistic to test for a trend in the ratio of ovarian:breast cancer risk along the gene, based on a chi-squared test for trend, as in (12) which was not

significant.

In an attempt to confirm the association using other data from previously published reports, we computed the chi-squared statistic (two degrees of freedom) using data from five studies (9,10,15-17), with the cutpoints n1 and n2 fixed at the best values for the UK dataset. The significance of this result was then based on permuting the mutations among families as before.

#### 10 Discussion

BRCA2 probably accounts for the majority of high risk families not accounted for by BRCA1. The identification of BRCA2 should, therefore allow more comprehensive evaluation of families at high risk of developing breast cancer. However, the role of environmental, lifestyle or genetic factors in modifying the risks of cancer in gene carriers is unknown and further studies will be required before routine diagnosis of carrier status can be considered.

20 Although, many of the mutations described in these examples are not is common to the BRCA2 genes of different families, these mutations may nevertheless prove to be a useful in methods of diagnosing or detecting a predisposition to cancer, especially breast cancer. In addition, on further routine analysis of a larger sample of families, it may emerge that a given mutation is common to a significant portion of cancer cases and so might form the basis of a simple diagnostic test. Even if this is not the case, knowing the location of the mutations may help to reduce the amount of sequencing work that would be needed in carrying out a test on a patient, e.g. by sequencing all or part of their BRCA2 gene, thereby to determine a predisposition or susceptibility to cancer. Further applications and uses of the mutations are set out in the general section above.

35 Germline mutations of BRCA2 are predicted to cause approximately 35% of families with multiple case, early onset female breast cancer, and they are also associated with an increased risk of male breast cancer, ovarian cancer, prostate cancer and pancreatic cancer (5,9,10). Germline mutations of a second cancer susceptibility gene BRCA1 (3), are associated with a strong predisposition to ovarian cancer as well as female breast cancer (11). Recent studies have suggested that the phenotype in BRCA1 families with respect to the ratio of breast to ovarian cancer varies with the location of the BRCA1 mutation (12,13). To determine whether germline mutations in BRCA2 are associated with a similar variation in phenotypic risk, we have analysed the distribution of mutations of families with multiple cases of breast and/or ovarian

5 cancer ascertained in the United Kingdom and Eire. The majority of these mutations lead to premature truncation of BRCA2 as a result of frameshift deletions, insertions, nonsense mutations or splice site alterations. Analysis of the mutation distribution along the length of the gene indicates a significant genotype-phenotype correlation. Truncating mutations in families with the highest risk of ovarian cancer relative to breast cancer are clustered in a region of approximately 3.3kb in exon 11 ( $p=0.0005$ ). Published data on mutations in 45 other BRCA2 linked families provide support for this correlation.

10 Families were ascertained on the basis of either three or more cases of breast cancer or two or more first degree relatives with epithelial ovarian cancer diagnosed at any age. Mutations of the BRCA1 gene were then identified and genomic DNA of a single affected individual from each of the remaining families was used to screen for mutations throughout the coding sequence of BRCA2 using a combination of single-strand conformation polymorphism (SSCP) and protein truncation (PTT) assays. The BRCA 2 coding sequence consists of 10,248 nucleotides encoded by 26 exons (15). The majority of exons are relatively small although exons 10 and 11 represent approximately 60% of the entire coding region. The mutations found are set out in tables 1 and 2. The mutation spectrum consisted of 22 frameshift deletions or insertions, 3 nonsense mutations and one splice site alteration. These mutations are all predicted to result in premature truncation of the predicted BRCA2 peptide. Three novel putative missense mutations were also detected. Mutations were distributed throughout the gene and only one additional mutation, 6503delTT, was found to be recurrent.

30 BRCA2 mutation data from families ascertained outside the UK derived from previous reports and unpublished data from our own laboratories provide support for this clustering. BRCA2 mutations have been identified in 45 such families (9,10,15-17). The 17 families with mutations in the OCCR are reported to contain 11 ovarian and 45 breast cancer cases compared to 22 ovarian and 282 breast cancer cases in the remaining families (odds ratio 22.9; permutation test for differences  $p=0.007$ ). Consistent with this, there are three large BRCA2 families where reasonably systematic evidence on cancer risks are available, namely UTAH 107 (15), CRC 186 and the Icelandic family (10). Mutations in all three families are located near the 5' or 3' end of the gene and appear to be associated with a high lifetime risk of breast cancer, with mutations in the OCCR being associated with a higher ovarian cancer risk than average, or a lower breast cancer risk, or both. In this regard, future data on the 6174delT mutation will be particularly important. This mutation, which lies in the OCCR, is common in Ashkenazi Jews and direct population based estimates of its prevalence

in breast and ovarian cancer patients should be possible.

The observed genotype-phenotype correlation is somewhat surprising given that, unlike in BRCA1, a region in the centre of the gene appears to be associated with a distinct risk. There are precedents in cancer susceptibility genes, for example the adenomatous polyposis coli gene, where regions of mutation clustering within the gene are associated with predisposition to a specific phenotype (18,19). We do not yet know why truncating mutations in a central portion of the BRCA2 gene should result in increased risk to ovarian cancer. No regions of functional homology between BRCA2 and other genes have been identified. A series of eight internal amino acid repeats have recently been observed in exon 11 but these have little homology to other known genes (20). It is perhaps interesting that all eight repeats are contained within the OCCR. That the mutations cluster in a single exon suggests an explanation based on alternative splicing. Complete or partial splicing of exon 11 may produce alternatively spliced forms with the ability to "rescue" mutations in breast but not ovarian epithelium. Our results suggest that they may be interesting differences in the structure or function of BRCA2 between breast or ovarian epithelium.

### Example 3 Expression of BRCA2 Polypeptide

#### Construction of Full-Length cDNA Clones for Human BRCA2

The approximately 10.5kb cDNA coding for human BRCA2 was assembled from fragments of cDNA amplified by PCR and from cloned genomic DNA. All residue numbers are from Genbank accession number U43746.

The 5' end of the cDNA (residues 204-2357) was amplified by PCR from cDNA made from RNA from MCF7 cells using the primers:

CATTGGAGGAATATCGTAGG (bases 204-223) and  
GACAGAGAATCAGCTTCTGG (bases 2338-2357).

This 2.1kb fragment was cloned into pBluescript and the sequence determined.

This plasmid (p12302) was then digested with Asp718 which cuts in the plasmid polylinker and partially with BamHI which cuts at bases 238 and 792. The double-stranded oligonucleotide:

GTACCGCCGCCATGGAACAGAAGATTCCGAAGAAGATCTGCCTATTG  
GCGGCGGTACCTTGTCTTCTAAAGGCTTCTTCTAGACGGATAACCTAG

was then ligated into the plasmid and recombinants carrying the oligo



ligated to the plasmid digested at base 238 were selected.

This plasmid was digested with NdeI which cuts at bp 1795 and SmaI which cuts in the polylinker of the plasmid. This was ligated in a three-way ligation to a PCR product (1795-2280) which had been digested with NdeI and BstY1 and residues 2280-6571 as a BstY1-BsiHKAI fragment. The latter fragment consists of exon 11 of the BRCA2 gene and was isolated from a plasmid carrying genomic DNA. This generated plasmid p12672.

The 3' end of the BRCA2 cDNA was generated by RT-PCR with primers:

AAAAGTAACGAACATTCAGACCA (bases 6277-6299) and  
ATTGTCGCCTTTGCAAATGC (bases 10,486-10,505) on cDNA from MCF7 cells.

This fragment was subcloned into pBluescript to generate plasmid p12661. Subsequently this fragment was amplified with a primer (GTACTCCAGAACATTTAATATCC bases 6325-6344) and the T3 primer from bluescript.

This fragment digested with Bsp1201 was then ligated into SnaBI-NotI cut p12672 to generate the full-length clone p12806.

This plasmid was subsequently modified to include a FLAG epitope as follows. p12806 was digested with Asp718 and SalI and then ligated to pBlueBac2B digested with the same enzymes. This generated plasmid p13013. This was digested with SacI and BgIII and ligated to a linker coding for a consensus Kozak sequence and a Flag epitope as detailed below:

CGGGTACCAGATCTGCCGCCACCATGGATTACAAGGACGACGATGACAAG  
TCGAGCCCATGGTCTAGACGGCGGTGGTACCTAATGTTCTGCTGCTACTGTTCTAG

This full-length BRCA2 cDNA was ligated into expression vectors. In particular a modified version of the vector pMTSM was used this vector carries an adenovirus major late promoter and SV40 origin of replication.

#### Expression of BRCA2 in Mammalian Cells

The full-length BRCA2 expression vector was transfected into COS cells using standard techniques. The expression of the protein was monitored by Western blotting immunoprecipitation and immunofluorescence. This demonstrated that the BRCA2 protein could be detected as an approximately 400kD protein similar in size to that predicted by the sequence of the cDNA. Under these conditions, in this cell-type, the

protein had a complex subcellular localisation being located in either a membrane-like compartment (most probably the endoplasmic reticulum) or in the nucleus or in both (see figures 11 to 13).

5      **Interaction of BRCA2 With Other Proteins**

This can be assessed using a yeast two-hybrid system to clone proteins which interact with BRCA2. As this procedure is potentially complicated by the size of the BRCA2 ORF, the ORF may be divided up into 5-10 fragments and screened separately. These fragments will be used as 'bait' both episomally and as integrants into the yeast genome and used to screen peptide libraries such as those derived from HeLa cells, human fetal brain and a new library derived from normal human breast. Clones confirmed as interacting in the two hybrid system will assayed for interaction both in vivo and in vitro. Bacterially expressed GEX-fusion proteins will be tested for direct interaction in vitro. Epitope-tagged versions of BRCA2 and the potential interacting proteins will be co-microinjected into cell lines and their sub-cellular location determined by immunofluorescence to establish co-localization. Co-transfection and cross-immunoprecipitation could be used to establish that the two proteins interact in vivo. In addition to identifying novel BRCA2 interacting proteins, the above approaches may be used to ascertain whether BRCA2 can dimerise or interact directly with BRCA1.

25      The nature of BRCA2 interacting proteins can also be directly determined by biochemical fractionation followed by mass spectroscopy. Immunoprecipitation of <sup>35</sup>S-labelled extracts followed by SDS-gel electrophoresis can be used produce molecular weight estimates of these proteins. These would be further analysed by analytical 2D gel electrophoresis followed by MALDI-TOF mass-spectroscopy and peptide mass-mapping (23). This technique allows the certain identification of proteins whose sequences are present in the databases and assignment of likely family members (>80% identity).

35      **Biological Function of BRCA2**

An experiment can be performed to test whether BRCA2 expression can block the growth of breast and ovarian cancer cell lines specifically. Ideally such experiments make use of breast tumour cells that do not express BRCA2, which can readily be identified by screening existing breast cancer cell lines for absence of BRCA2 expression. Alternatively, cell lines could be established from patients shown to lack wild type BRCA2. It is also possible that over-expression of BRCA2 in cancer cells that still express will also suppress growth.

45      BRCA2 cDNAs can suitably be expressed under the control retroviral LTR

(pBABE) or elongation factor 1 $\alpha$  promoters (the pEFBos series - (22)). Plasmids can be co-transfected with drug resistance markers and the number of colonies that grow out compared to vector controls. surviving colonies can be expanded and tested for tumourgenecity by injection into nude mice. In instances where the inhibitory action of the proteins cannot be detected in the relatively long term colony growth assay as the transfected plasmids do not stably express, microinjection of BRCA2 expression constructs can be used whereby the injected cells are detected by immunocytochemistry of the exogenous protein or by co-injection of a marker plasmid (21). In addition, time-lapse video recording, as described above, could be used to determine whether any growth inhibition effect is cell autonomous, i.e. whether the effect is paracrine or autocrine. These systems will also be useful for detecting any apoptotic effect.

#### Example 5    Methods for Detecting BRCA2 Mutations

Migration shift assays to detect BRCA2 mutations.

#### DNA amplification in the PCR.

25ng of genomic DNA from each individual to be screened for mutations is amplified in 35 cycles of the PCR using the pertinent oligonucleotide primers (see primer list). Prior to incorporation into the PCR, both oligonucleotide primers are end radiolabelled with gamma <sup>32</sup>p using T4 polynucleotide kinase. Following amplification in the PCR, formamide loading dye is added to each sample and the sample denatured at 94°C for 3 minutes. Following denaturation the sample is placed immediately on ice.

#### DNA fragment sizing.

2 $\mu$ l of each sample is loaded immediately onto a well formed by a 40 slot sharks' tooth comb in conventional 0.4mm thick denaturing 6% polyacrylamide gel. The sample is electrophoresed through the gel for 2-5 hours at 90 Watts at room temperature.

#### SSCP heteroduplex analysis

SSCP is a PCR based assay for screening DNA fragments for sequence variants/mutations. It involves amplifying radiolabelled 100-300 bp fragments of the BRCA2 gene, diluting these products and denaturing at 95°C. The fragments are quick-cooled on ice so that the DNA remains in single stranded form. These single stranded fragments of BRCA2 are run through acrylamide based gels. Differences in the sequence composition will cause the single stranded molecules to adopt difference conformations in this gel matrix making their mobility different from wild type fragments, thus allowing detecting of mutations in the

fragments being analysed relative to a control fragment upon exposure of the gel to X-ray film.

5 These fragments with altered mobility/conformations are directly excised from the gel and directly sequenced for the mutation. Following denaturation the sample is cooled on ice for 10 minutes to allow the heteroduplex to form. Each sample is electrophoresed through two different types of gel.

10 A typical set of conditions for SSCP analysis are as follows. 3 $\mu$ l are electrophoresed overnight at 4 Watts at room temperature through a 6% non denaturing polyacrylamide gel containing 10% glycerol.

15 3 $\mu$ l are electrophoresed for four hours at 30 Watts in a 4°C cold room through a 4.5% non denaturing polyacrylamide acrylamide gel without glycerol.

20 Following electrophoresis, gels are dried onto Whatman 3MM paper, and placed in an autoradiography cassette at room temperature for a period ranging from two hours to several days.

Following development of the autoradiograph band shifts in individual samples are detected by eye.

#### 25 Sequencing of PCR product.

Where a band shift is seen in SSCP heteroduplex or DNA fragment sizing gels, the fragment concerned is reamplified from the relevant stock genomic DNA and directly sequenced. To sequence PCR product, the product is precipitated with isopropanol, resuspended and sequenced using TaqFS+ Dye terminator sequencing kit. Extension products are electrophoresed on an ABI 377 DNA sequencer and data analysed using Sequence Navigator software.

#### BRCA2 PTT Assay

35 PTT is another PCR based screening assay. Fragments of BRCA2 are amplified with primers that contain the consensus Kozak initiation sequences and a T7 RNA polymerase promoter. These extra sequences are incorporated into the 5' primer such that they are in frame with the native coding sequence of the fragment being analysed. These PCR products are introduced into a coupled transcription/translation system. This reaction allows the production of BRCA2 RNA from the fragment and translation of this RNA into a BRCA2 protein fragment. PCR products from controls make a protein product of a wild type size relative to the size of the fragment being analysed. If the PCR product analysed has a frame-shift or nonsense mutation, the assay will

45

yield a truncated protein product relative to controls. The size of the truncated product is related to the position of the mutation, and the relative region of the BRCA2 gene from this patient is sequenced to identify the truncating mutation.

5

The following protocol was adapted for a BRCA2 PTT assay. The PTT primers are shown in figure 10.

Each PTT primer is preceded by the T7/Kozak sequence  
10 GGATCCTAATACGACTCACTATAGGGAGACCACCATG

1. Thirty five cycle primary PCR reaction in 20ul.
2. Product confirmation by electrophoresis on 2% agarose gel.
3. Three  $\mu$ l aliquot amplification with nested PTT prime for 15  
15 cycles.
4. Product confirmation by electrophoresis on 2% agarose gel.
5. In vitro transcription/translation using Promega (CA) TNT kit, incorporating  $^{35}$ S radiolabelled methionine.
6. Laemmli buffer reaction stop and denaturation.
- 20 7. Gel electrophoresis of product on 15% acrylamide gel at 16mA.
8. Fix, amplify and dry gel.
9. Autoradiographic exposure for 2 hours.

25 These approaches can be combined to provide an accurate and effective screen, in terms of results achieved, the economical cost and the time taken to provide the results. By way of example, a combined protocol used by the inventors involves the following:

30 1) DNA samples from familial reference cases (probands) are first screened via PTT. Exons 10, 11, and the terminal exon 27 are analysed using this technique. Exons 10 and 27 are done in 1 fragment, whilst the larger exon 11 is done in 2 fragments. If protein truncations are seen, the corresponding genomic region is sequenced from the patient to identify the exact mutation. This approach is able to screen  
35 approximately 60% of the coding region in rapid fashion.

2) Samples negative for the PTT screening are then analysed using SSCP. The entire coding region, including those regions already examined via PTT, is screened for mutations using SSCP. The coding  
40 sequence is amplified from genomic DNA using primer sets as described in this application. In addition, radiolabelled PCR products generated for SSCP analysis are also run on denaturing acrylamide sequencing gels which allows for detection of small size changes relative to control fragments, indicative of insertions or deletions. Any mutant fragments  
45 seen are excised directly from the gels and direct sequenced along with

the matching genomic DNA fragment to determine the exact mutation

Examples of screening carried out using these methods are set out in the table below.

5

BRCA2 screening in breast cancer family probands:

Family	Mutation	No. Female Br Ca	Exon	Method of Detection
10 M107	2810insA	8	11	SSCP/PTT, mutation detected on both assays
M742	259insT	6	2	SSCP
M688	594delCT	2 (2 Ov)	11	PTT
15 M347	1003delAG	3	9	SSCP
D-MBC2	1485delT	5 (1 Mbr)	10	PTT
M269	8764delAG	19	20	SSCP
M612	8764delAG	4	20	SSCP
20 M133	2034insA	11	10	SSCP, too close to end of exon to be reliably up by PTT

Example 6 Results of Sequence Alignment

25 Materials and Methods

Total genomic DNA was obtained for green monkey (*Ceropithecus aethiops*), hamster (*Critetulus griseus*), pig (*Sus scrofa*), dog (*Canis familiaris*), cow, chimpanzee, chicken, snake, zebra fish, *X.laevis*, *S.pombe* and *S.cerevisiae*. PCR primers were designed from a consensus sequence for the human BRC motif. These included primers that were an exact match to the human DNA sequence and degenerate primers based on the BRCA2 protein sequence. The sequences were as follows.

35 Motif forward set 1: pure primer -AAAGCTGTGAAACTGTT and a degenerate set which was an equal mixture of -AA(AG)GCIIIIIAA(AG)CTITT and -AA(AG)GCIIIIIAA(AG)TT(AG)TT where I is inosine.

Motif forward set 2: pure primer as for set 1 and a degenerate set made with -AA(AG)GCIGTIAA(AG)CTITT and -AA(AG)GCIGTIAA(AG)TT(AG)TT.

40 Motif reverse set: pure primer - TTCCCACTTGCACTCTGAAA and a mixed set of -TICC(GA)CTIGCIGT(CT)TG(GA)AA and -TTICCIGAIGCIGT(CT)TG(GA)AA.

45 Further sets of primers were designed to a region of BRCA2 that is conserved between human and mouse. These were a pure primer of

AGCAAGCAATTTCAAGG and a degenerate set of -  
TCIAA(AG)CA(AG)TT(TC)GA(AG)GG and -AG(TC)AA(AG)CA(AG)TT(TC)GA(AG)GG.  
The PCR was performed using each forward primer in conjunction with a  
relevant reverse primer. The reaction conditions were as described in  
5 (4) except for the cycling conditions of the PCR. These were 94°C 1 min  
X°C for 1 min. 72°C for 1 min, where X was 65°C for the first two  
cycles and then decreased by 2°C every two cycles until it reached  
either 55, 49, 45, 39 or 35°C, at which point X remained constant. A  
further 30 cycles were performed at the final annealing temperature.  
10 PCR products were resolved on agarose gels and discrete bands were  
excised and sequenced using a Taq dideoxy termination protocol (Perkin-  
Elmer). Sequence products were separated on an ABI377 automated DNA  
sequencer.

15 All of the sequencing reactions were performed at least twice and on  
both strands of the template. Species-specific PCR primers were  
designed to exon 11 of the BRCA2 gene using the above DNA sequence.  
These were used in conjunction with a set of 52 pairs of PCR primers  
designed to the human BRCA2 gene to amplify and sequence further  
20 segments of exon 11 from the monkey, dog, pig and hamster. A human  
BRCA2 probe (exon 11) was used to identify a  $\lambda$  clone containing a  
portion of the mouse BRCA2 gene. This was used to obtain a mouse-  
specific BRCA2 sequence that was in turn used to design mouse-specific  
PCR primers. These were used to identify positive clones in a mouse  
25 BAC library. Fragments of positive BACs were cloned, at random, and  
those shown to contain fragments of exon 11 of the BRCA2 gene were  
sequenced.

#### Sequencing the BRC motifs

30 Two approaches were used to obtain the sequence of BRCA2 exon 11  
(including the BRC motifs) in five mammalian species, using the  
sequence of human BRCA2 was taken from earlier work, see also figure 7.  
The sequence of exon 11 of the mouse BRCA2 gene was obtained from a BAC  
clone isolated by low stringency hybridisation with fragments of the  
35 human BRCA2 gene. Fragments of BAC were cloned at random and those  
shown to contain fragments of the BRCA2 gene were sequenced. PCR  
primers were then designed to the BRC motifs and to regions that were  
conserved between human and mouse. These were used to amplify  
fragments of the BRCA2 gene in DNAs from 12 species (as described in  
40 Materials and Methods). A sequence showing similarity to human BRCA2  
was obtained from green monkey, hamster, pig, dog, cow and chimpanzee.  
The sequences obtained from chicken, snake, zebra fish, *Xenopus laevis*,  
*Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* showed no  
similarity to any BRCA2 sequence. Species-specific PCR primers were  
45 designed for green monkey, hamster, pig and dog. These were used in

combination with all of the PCR primer pairs used to amplify human BRCA2 exon 11 to amplify further segments of the exon. The process was repeated until all of the sequences between and including BRC motifs one and eight had been obtained for the green monkey, hamster and dog. Using this approach, it was only possible to obtain the sequence for repeats 3-8 for the pig. In total, 46 BRC repeats were sequenced in six different mammals (including human).

**The BRC motif is conserved in mammalian species**

The percentage identity of the translation of exon 11 of the BRCA2 gene (approximately half of the whole coding sequence) between the six mammals is shown in Table 1. Overall, the degree of conservation is low, with 58% identity between the 1602 residues of the human and mouse, 54% identity between mouse or hamster and dog and 49% between pig and mouse (over 928 amino acids). Even between closely related species such as human and monkey, and mouse and hamster, amino acid identity is only 93 and 72% respectively. However, an alignment of the sequences from the six mammals studied demonstrates that exon 11 translations can be aligned along their whole lengths and that the degree of conservation is variable (figure 9). There are a number of short regions of high identity. Some of these regions coincide with BRC motifs, for example BRC1, 2, 3, 4, 7 and 8 (figure 9). There are other highly conserved segments, for example amino acids 469-493 (using the numbering of the human sequence in figure 9) that show no similarity to the BRC motifs or anything else reported in the databases. None of the latter are repeated within BRCA2.

An alignment of all 46 motifs sequenced demonstrates the high degree of interspecies and intraspecies conservation between BRC1, 3, 4, 7 and 8. From this alignment, we have identified a region of 26 amino acids that is conserved in all of the BRC motifs (Fig. 2) which has allowed us to generate a BRC consensus sequence. It is possible to align some residues outside this common region; however, such alignments are not robust, being very sensitive to the parameters used. There are motifs that contain all of the consensus sequence, for example BRC4, while others, for example BRC6, show a considerable divergence from the consensus. In total, 30/46 of the BRC motifs (65%) have 11 or more of the 13 consensus residues while 87% have eight or more of the consensus sequence.

The following table shows the percentage identities between the translation of exon 11 of the BRCA2 gene from human, monkey, pig, dog, hamster and mouse.



	Human	Monkey	Pig	Dog	Hamster	Mouse
Human	100	93	62	70	57	58
Monkey		100	61	68	58	58
Pig			100	64	50	49
Dog				100	54	54
Hamster					100	72
Mouse						100

### Discussion

The low sequence identities for exon 11 (shown in the table above) suggest that BRCA2 is evolving at a faster rate than most other cancer susceptibility/tumour suppressor genes. For example, there is 98% identity between human and mouse NF1 with 95 and 91% between human and mouse WT1 and Rb1 respectively. A notable exception, however, is BRCA1 which only shows 58% amino acid identity between mouse and human. Therefore, although BRCA1 and BRCA2 do not show substantial sequence similarity, they are similar by virtue of a high rate of evolution, unusual gene structure (both have a large exon 11, see (3) and (7) and lack of somatic mutations in sporadic cancers. Whether one or more of these features relates to a function congruence remains to be elucidated.

Although the identity for the translation of exon 11 of the BRCA2 gene is low, most of the BRC motifs are highly conserved between the species analysed. This suggests that there has been pressure to maintain the BRC repeats in BRCA2 and, therefore, that they are important in its function. However, from our alignment, BRC6 is much less conserved than BRC1, 2, 3, 4, 7 and 8 (31 and 85% identity between human and mouse BRC6 and BRC7 respectively). Moreover, it has been altered by insertions or deletions such that the length of the motif differs between species. BRC3 and BRC5 also shows less conservation than BRC1, 2, 3, 4, 7 and 8 (62 and 58% identity respectively between human and mouse BRC3 and BRC5). However, both exhibit a higher level of conservation than the exon 11 translation overall. The data suggests that multiplication of the motif took place prior to the mammalian radiation. For instance, several amino acid residues within motif units (especially in BRC2) are different from the equivalent residues in other units, but are highly conserved in mammalian species. Moreover, the sequences flanking the motifs are not conserved between repeats, but in some cases are conserved across species (for example BRC1 and 4, figure 9).

The PCR products that we derived from non-mammalian DNA did not exhibit any similarity to mammalian BRCA2. This suggests that either BRCA2 is restricted to mammals or that the non-mammalian orthologues of BRCA2 have diverged to such an extent that they cannot be identified by the techniques that we used. However, there is evidence of a weak similarity between the BRC sequence and a *Caenorhabditis elegans* gene (CET07E3\_2), suggesting that the BRC motif is not restricted to mammals.

There are few clues to the function of the BRC sequences. Truncating germline mutations in the BRCA2 gene that predispose to the development of breast cancer are located throughout the coding region of the gene. In many cases, the mutations leave all the BRC repeats intact, giving little information on the relationship between the motifs and their role in the normal function of BRCA2. The spacing between individual motifs varies from -60 to 300 amino acids, but is reasonably well conserved between mammals. Furthermore, there are multiple elements of secondary structure within each motif which may indicate that they form globular domains. However, it is unclear how these may function. Therefore, direct investigation will be necessary to elucidate the biochemical functions of the BRC motifs.

Conserved regions such as the BRC motifs identified in exon 11 provide valuable indications of domains of the BRCA2 polypeptide that are likely to be important in determining its activity. Thus, these motifs are good candidates for screening studies described above, to find mimetics for BRCA2 polypeptide.

Motifs from exon 11 that are conserved between some or all of the species examined and depicted in figure 9 and are summarised in table 4:

**Table 4**

	<u>Start Amino Acid</u>	<u>End Amino Acid</u>	<u>Motif</u>
35	379	404	SFRTASNKEIKLSEHNIKKSKMFFKD
	594	619	GFYSALGTKLNVSEALQKAVKLFSD
	806	831	SFQTASGKNIRVSKESLNKAVNFFDQ
	909	934	GFHTASGKKVKIAKESLDKVKNLDFE
	1057	1082	AFYTGHSRKTSVSEASLLEAKKWLRE
40	1257	1271	AFSTASGKIVFVSHE
	1392	1417	IFSTASGKSVQVSDASLQKARQVFSE
	1474	1499	GFSTASGKQVSVSESALHKVKGMLEE

Table 1. Germline mutations of the BRCA2 gene in breast and ovarian cancer families

5	Family	Number of cases of cancer		Description of mutation				
		Breast	Breast ≤ 60 yrs	Ovarian	Exon	Codon	Nucleotide Alteration	Mutation Type#
10	B061	2	2	0	2	9	253delC	FS
	B091	4	4	0	7	194	809 G>T	N
	B135	3	3	0	10	397	1418	FS
	B105	2	2	0	10	434	1529	FS
	B022	4	3	0	10	598	2022del5	FS
15	B169	3	3	0	11	643	2157delG	FS
	B124	3	3	0	11	936	3034del4	FS
	B211	4	4	4	11	2092	6503delTT	FS
	B150	6	5	0	11	2092	6503delTT	FS
	B160	4	2	0	11	2134	6630del5	FS
20	B245	7	3	0	11	2154	6690delTC	FS
	B196	18	14	0	11	2197	6819delTG	FS
	B097	4	3	0	11	2277	7057del5	FS
	B007 <sup>d</sup>	11	5	0	17	2637	8138del5	FS
	B028 <sup>d</sup>	3	3	0	17	2654	8162delG	FS
25	B186	18	18	1	22	2984	9179C>G	N
	B247	3	3	0	23	2994	9208ins4	FS
	B001	4	3	0	25	3167	9729G>A	SS
	M681 <sup>d</sup>	3	3	0	11	1982	6174delT	FS
	M440 <sup>d</sup>	2	2	0	11	936	3034del4	FS

<sup>d</sup>These families also contain one or two male breast cancer cases. #Mutation types characterised as frameshift deletions or insertion (FS), nonsense mutations (N), missense mutations (M) or splice site alterations (SS).

Table 1 (cont). Germline mutations of the BRCA2 gene in breast and ovarian cancer families

	5	Family	Number of cases of cancer			Description of mutation			
			Breast	Breast ≤ 50 yrs	Ovarian	Exon	Nucleotide	Alteration	Mutation Type#
10		M107 8	6	0		11	2810	2810insa	FS
		M133 11	4	1		10	2034	2034insa	FS
		M174 9	8	0		11	3034	3034del4	FS
		M269 <sup>d</sup> 21	11	0		20	8764	8764delAG	FS
		M280 4	0	0		11	6085	G6085Tns	N
15		M284 3	2	1		11	4304	4304delC	FS
		M347 <sup>d</sup> 4	0	0		9	1003	1003delAG	FS
		M372 6	0	0		25	9661	9661delGT	FS
		M609 12 <sup>d</sup>	0	0		10	2024	2024del5	FS
		M612 4	3	0		20	8764	8764delAG	FS
20		M688 2	0	2		11	5946	5946delCT	FS
		M733 2	0	0		11	6437	6437del4	FS
		M742 6	4	0		2	259	259inst	FS
		W9052 4	3	0		10	2034	2034delA	FS
		W9116 4	3	0		10	1917	G1917Ans	N
25		DWBC2 6 <sup>d</sup>	2	0		10	1485	1485delT	FS
		W9012 1 <sup>d</sup>	0	0		11	5839	5839del4	FS

<sup>d</sup>These families also contain one or two male breast cancer cases. #Mutation types characterised as frameshift deletions or insertion (FS), nonsense mutations (N), missense mutations (M) or splice site alterations (SS).

TABLE 2

Family	Mutation description	Mutation type	sub sequence	d/i sequence	Location exon	Location fragment	Location base	Location codon
5								
CRC B001	Splice	s	G->A		25	B	9729	0
CRC B007	Frameshift	d5		CCTTT	17		8138	2637
CRC B022	Frameshift	d5		ATCTT	10	E	2022	598
10								
CRC B028	Frameshift	d1		G	17		8162	2645
CRC B061	Frameshift	d1		C	2		253	9
CRC B097	Frameshift	d5		CTTAT	11	Z	7057	2277
CRC B105	Frameshift	d4		AAAG	10	B	1529	434
CRC B124	Frameshift	d4		AAAC	11	F	3034	936
15								
CRC B135	Frameshift	i4		TTAG	10	B	1418	397
CRC B150	Frameshift	d2		TT	11	W	6503	2092
CRC B160	Frameshift	d5		TAACT	11	X	6630	2134
CRC B169	Frameshift	d1		G	11	A	2157	643
CRC B196	Frameshift	d2		TG	11	Y	6819	2197
20								
CRC B211	Frameshift	d2		TT	11	W	6503	2092
CRC B245	Frameshift	d2		TC	11	X	6690	2154
CRC B247	Frameshift	i4		AGAT	23		9208	2994
IARC3594	Frameshift	d2		CT	11		5946	1906
IARC2932	Frameshift	d6		TTTCGg	13		7231	2334
25								

TABLE 3

	Number of FBCs	Number of FBCs <50	Number of OVCs	Number of MBCs	LOD score at BRCA1	LOD score at BRCA2	BRCA2 mutation
IARC 2932	15	10	0	0	-2.38	3.01	CCC.TTT.CGgtaa
IARC 3594	6	5	0	0	nd	nd	CAT.AAC.TCT.CTA
CRC B211	5	3	4	0	-0.48	0.49	AGT.CTT.CAC
CRC B196	17	12	0	0	-2.21	0.92	AAA.ACT.GAA.ACT
Montreal 681	3	2	0	1	nd	nd	GCA.AGT.GGA
Montreal 440	2	2	0	2	nd	nc	GAT.AAA.CAA.GCA

References:

The references cited below and in the above description are all incorporated by reference in their entirety.

- 5 1. Hall et al. Linkage of early-onset familial breast cancer to chromosome 17q21. *Science*, 250(4988):1684-1689, 1990.
2. Malkin et al. Germ line p53 mutations in a familial syndrome of  
10 breast cancer, sarcomas, and other neoplasms. *Science*, 250(4985):1233-1238, 1990.
3. Miki et al. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science*, 266(5182):66-71, 1994.  
15
4. Wooster et al. A germline mutation in the androgen receptor gene in two brothers with breast cancer and Reifenstein syndrome. *Nat. Genet.*, 2(2):132-134, 1992.
- 20 5. Wooster et al. Localization of a breast cancer susceptibility gene, BRCA2, to chromosome 13q12-13. *Science*, 265(5181):2088-2090, 1994.
6. Frohman et al. Rapid amplification of full length cDNAs from rare transcripts: Amplification using a single gene-specific oligonucleotide  
25 primer. *PNAS USA*, 85:8998-9002, 1988.
7. Wooster et al. Identification Of the Breast-Cancer Susceptibility Gene BRCA2. *Nature*, 378:789-792, 1995.
- 30 8. Beaudet and Tsui. A Suggested Nomenclature for Designating Mutations. *Hum. Mutat.*, 2(4):245-8, (1993).
9. Thorlacius et al. A Single BRCA2 Mutation In Male and Female Breast-Cancer Families From Iceland With Varied Cancer Phenotypes. *Nature  
35 Genetics*, 13:117-119, 1996.
10. Phelan et al. Mutation Analysis Of the BRCA2 Gene In 49 Site-Specific Breast-Cancer Families. *Nature Genetics*, 13:120-122, 1996.
- 40 11. Easton et al. Genetic Linkage Analysis in Familial Breast and Ovarian Cancer. *Am. J. Hum. Genet.*, 52:718-722, 1993.
- 45 12. Gayther et al. Germline Mutations Of the Brcal Gene In Breast and Ovarian-Cancer Families Provide Evidence For a Genotype-Phenotype Correlation. *Nature Genetics*, 11:428-433, 1995.

13. Holt et al. Growth-Retardation and Tumor-Inhibition by BRCA1. Nature Genetics, 12:298-302, 1996.
14. Gayther et al. Rapid Detection Of Regionally Clustered Germ-Line BRCA1 Mutations By Multiplex Heteroduplex Analysis. Am. J. Human Gen., 58:451-456, 1996.
15. Tavtigian et al. The Complete BRCA2 Gene and Mutations In Chromosome 13q-Linked Kindreds. Nature Genetics, 12:333-337, 1996.
16. Couch et al. BRCA2 Germline Mutations In Male Breast-Cancer Cases and Breast-Cancer Families. Nature Genetics, 13:123-125, 1996.
17. Neuhausen et al. Recurrent BRCA2 6174delT Mutations In Ashkenazi Jewish Women Affected By Breast-Cancer. Nature Genetics, 13:126-128, 1996.
18. Spirio et al. Alleles of the APC gene: an attenuated form of familial polyposis. Cell, 75:951-957, 1993.
19. Olschwang et al. Restriction of ocular fundus lesions to a specific subgroup of APC mutations in adenomatous polyposis coli patients. Cell, 75:959-968, 1993.
20. Bork et al. Internal Repeats In the BRCA2 Protein-Sequence. Nature Genetics, 13:22-23, 1996.
21. Cowley, S., Paterson, H.F., Kemp, P. and Marshall, C.J., Cell, 77:841-852, 1994.
22. Marais, R., Light, Y., Paterson, H.F. and Marshall, C.J., EMBO J., 14:3136-3145, 1995.
23. Hynes, G. et al, FASEB J. 10:137-147, 1996.



Claims:

1. A nucleic acid molecule comprising a part of the BRCA2 gene as set out in figures 1 or 2, or alleles thereof.
- 5 2. A nucleic acid molecule comprising the full length coding sequence or complete BRCA2 gene as obtainable by:
- (a) using the nucleic acid sequences shown in figures 1 or 2 to construct probes for screening cDNA or genomic libraries, sequencing the positive clones obtained, and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained;
- 10 (b) using the sequences shown in figures 1 or 2 to obtain oligonucleotides for priming BRCA2 nucleic acid fragments, these oligonucleotides being used in conjunction with oligonucleotides designed to prime from a cloning vector, to amplify by PCR nucleic acid fragments in a library that contains fragments of the BRCA2 sequence, sequencing the amplified fragments to obtain the BRCA2 sequence between known parts of the sequence and the cloning vector, and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained; and/or,
- 15 (c) using rapid amplification of cDNA ends (RACE), by synthesizing cDNAs from a number of different RNAs, the cDNAs being ligated to an oligonucleotide linker, and amplifying by PCR the BRCA2 cDNAs using one primer that primes from the BRCA2 cDNA sequence of figure 1 and a second primer that primes from the oligonucleotide linker, sequencing the amplified nucleic acid and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained.
- 20 3. A nucleic acid molecule comprising a part of the BRCA2 gene as set out in figure 4 or alleles thereof.
- 30 4. A nucleic acid molecule comprising the full length coding sequence or complete BRCA2 gene as obtainable by:
- (a) using the nucleic acid sequences shown in figures 1, 2 or 4 to construct probes for screening cDNA or genomic libraries, sequencing the positive clones obtained, and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained;
- 35 (b) using the sequences shown in figures 1, 2 or 4 to obtain oligonucleotides for priming BRCA2 nucleic acid fragments, these oligonucleotides being used in conjunction with oligonucleotides designed to prime from a cloning vector, to amplify by PCR nucleic acid fragments in a library that contains fragments of the BRCA2 sequence, sequencing the amplified fragments to obtain the BRCA2 sequence between known parts of the sequence and the cloning vector, and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained; and/or,
- 40 (c) using rapid amplification of cDNA ends (RACE), by synthesizing
- 45

cdNAs from a number of different RNAs, the cdNAs being ligated to an oligonucleotide linker, and amplifying by PCR the BRCA2 cdNAs using one primer that primes from the BRCA2 cdNA sequence of figures 1 or 4 and a second primer that primes from the oligonucleotide linker, sequencing the amplified nucleic acid and repeating this process to assemble the full length BRCA2 sequence from the sequences thus obtained.

- 5 5. A nucleic acid molecule which is an allele or variant of a BRCA2 nucleic acid molecule as obtainable in claim 2 or claim 4.
- 10 6. A nucleic acid molecule which has a nucleotide sequence encoding a BRCA2 polypeptide including the amino acid sequence set out in figure 3.
7. A nucleic acid molecule which has a nucleotide sequence encoding a BRCA2 polypeptide including the amino acid sequence set out in figure 5.
- 15 8. A nucleic acid molecule which has a nucleotide sequence encoding a BRCA2 polypeptide including the amino acid sequence set out in figure 7.
- 20 9. A nucleic acid molecule which has a nucleotide sequence encoding a polypeptide which is a variant, derivative or allele of a BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5 or 7.
- 25 10. The nucleic acid of claim 9 encoding a polypeptide having at least 80% sequence homology to the BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5 or 7.
11. The nucleic acid molecule of claim 9 or claim 10 wherein the molecule has one or more of the mutations set out in table 1.
- 30 12. The nucleic acid molecule of claim 11 having a 6174delT mutation.
13. The nucleic acid molecule of claim 11 having a 6503delTT mutation.
- 35 14. A nucleic acid molecule which has a nucleotide sequence encoding a fragment or active portion of a BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5, or 7.
15. The nucleic acid molecule of any one of claims 1 to 14 further comprising all or a part the BRCA2 promoter region, the nucleic acid sequence of which is set out in figure 6.
- 40 16. A replicable vector comprising nucleic acid of any one of claims 1 to 14 operably linked to control sequences to direct its expression.

17. The vector of claim 16 further comprising the nucleic acid of claim 15 operably linked to promote the expression of the nucleic acid encoding the BRCA2 polypeptide.
- 5 18. Host cells transformed with the vector of claim 16 or claim 17.
19. A method of producing a BRCA2 polypeptide comprising culturing the host cells of claim 18 so that BRCA2 polypeptide is produced.
- 10 20. The method of claim 19 comprising the further step of recovering the polypeptide produced.
21. The nucleic acid molecule of any one of claims 1 to 15, or its complement, further comprising a label.
- 15 22. A nucleic acid molecule of any one of claims 1 to 15 for use in a method of medical treatment.
23. A substance which is a BRCA2 polypeptide encoded by the nucleic acid of claim 2 or claim 4.
- 20 24. A substance which is a BRCA2 polypeptide including the amino acid sequence set out in figure 3.
- 25 25. A substance which is a BRCA2 polypeptide including the amino acid sequence set out in figure 5.
26. A substance which is a BRCA2 polypeptide including the amino acid sequence set out in figure 7.
- 30 27. A substance which is a polypeptide having at least 80% sequence homology to the BRCA2 polypeptide including the amino acid sequence set out in any one of figure 3, 5 or 7.
- 35 28. A substance which is a polypeptide which is a variant, derivative or allele of a BRCA2 polypeptide of any one of claims 22 to 27.
29. A substance which is a fragment or active portion or functional mimetic of a BRCA2 polypeptide including the amino acid sequence of any one of figures 3, 5, or 7.
- 40 30. The substance of any one of claims 22 to 29 further comprising a label.
- 45 31. The substance of any one of claims 22 to 30 for use in a method of

medical treatment.

32. An antibody capable of specifically binding to a BRCA2 polypeptide of any one of claims 22 to 30.

33. The antibody of claim 32 further comprising a label.

34. A pharmaceutical composition comprising a substance of any one of claims 22 to 30.

35. A pharmaceutical composition comprising an antibody of claim 32 or claim 33.

36. The pharmaceutical composition of claim 34 or claim 35 additionally comprising a pharmaceutically acceptable carrier.

37. A method of diagnosing a susceptibility or predisposition to cancer in a patient by analysing a sample from the patient for the BRCA2 gene or the polypeptide encoded by it, the method comprising:

(a) comparing the sequence of nucleic acid in the sample with the BRCA2 nucleic acid sequence as set out in figure 1, 2, 4 or 7 to determine whether the sample from the patient contains mutations; or,

(b) determining the presence in the sample of the polypeptide encoded by the BRCA2 gene as set out in the partial sequences of figures 3 and 5 or the full length sequence of figure 7 and, if present, determining whether the polypeptide is full length, and/or is mutated, and/or is expressed at the normal level; or,

(c) using DNA fingerprinting to compare the restriction pattern produced when a restriction enzyme cuts a sample of nucleic acid from the patient with the restriction pattern obtained from normal BRCA2 gene comprising the sequence set out in figures 1, 2, 4 or 7 or from known mutations thereof; or,

(d) using a specific binding member capable of binding to a BRCA2 nucleic acid sequence as set out in figures 1, 2, 4 or 7 (either a normal sequence or a known mutated sequence), the specific binding member comprising nucleic acid hybridisable with the BRCA2 sequence, or substances comprising an antibody domain with specificity for a native or mutated BRCA2 nucleic acid sequence or the polypeptide encoded by it, and detecting the binding of the specific binding member to its binding partner by means of a label; or,

(e) using PCR involving one or more primers based on normal BRCA2 gene sequence set out in figures 1, 2, 4 or 7, or mutated forms thereof, to screen for normal or mutant BRCA2 gene in a sample from a patient.

38. The method of claim 37 wherein the patient sample is analysed by

single stranded conformation polymorphism (SSCP) and/or by a protein truncation (PTT) assay.

39. A method of identifying a target nucleic acid molecule in a test  
5 sample using a nucleic acid probe having a portion of the sequence shown  
in any one of figures 1, 2, 4 or 7 or a complementary sequence thereof, the  
method comprising contacting the probe and the test sample under  
hybridising conditions and observing whether hybridisation takes place.
- 10 40. The method of claim 39 wherein the probe is used to identify a BRCA2  
nucleic acid sequence or a mutant allele thereof.
41. The method of claim 39 wherein the probe is used to identify BRCA2  
nucleic acid of other species.
- 15 42. A method of determining the presence of one or more mutations in a  
sample of nucleic acid from a patient using one or more allele specific  
nucleic acid probes having the sequence or a portion of the sequence set  
out of figures 1, 2, 4 or 7 or a complementary sequence thereof, the method  
20 comprising contacting the probe and the test sample under hybridising  
conditions and observing whether hybridisation takes place.
43. The method of claim 42 wherein a plurality of allele specific probes  
are used, the probes being immobilised in an array on a solid support, the  
25 hybridisation of the nucleic acid sample to the probes being detected by  
means of a label.
44. The method of claim 42 or claim 43 wherein the mutations include one  
or more of those set out in table 1.
- 30 45. The use of a nucleic acid molecule of any one of claims 1 to 15 in  
the preparation of a medicament for treating cancer.
46. The use of claim 45 wherein the cancer is female breast cancer, male  
35 breast cancer, ovarian cancer, prostate cancer, colorectal cancer, ocular  
melanoma or leukaemia.
47. The use of claim 45 or claim 46 wherein the nucleic acid molecule  
is an antisense oligonucleotide capable of hybridising to the complementary  
40 sequence of BRCA2 nucleic acid, pre-mRNA or mature mRNA so that the  
expression of the BRCA2 nucleic acid is reduced or prevented.
48. The use of claim 45 or claim 46 wherein the use of the nucleic acid  
is in a method of gene therapy.
- 45

49. The use of a nucleic acid sequence of any one of claims 1 to 15 in the design of primers for use in the polymerase chain reaction.
50. The use of a nucleic acid sequence of any one of claims 1 to 15 in the design of a nucleic acid probe for detecting the presence of mutations in a nucleic acid sample from a patient.
51. The use of nucleic acid encoding all or a functional part of the BRCA2 promoter region set out in figure 6 in screening for substances which modulate the expression of nucleic acid under the control of the promoter.
52. A polypeptide having a BRCA2 amino acid motif set out in table 4, the motif being conserved between the human BRCA2 sequence set out in any one of figures 3, 5 or 7 and the sequence of another species.
53. Use of a substance of any one of claims 23 to 30 in the preparation of a medicament for treating cancer.
54. The use of claim 53 wherein the cancer is female breast cancer, male breast cancer, ovarian cancer, prostate cancer, colorectal cancer, ocular melanoma or leukaemia.
55. The use of an antibody of claim 32 or claim 33 for determining the presence, amount or location in a cell of BRCA2 polypeptide or mutant forms thereof.
56. Use of substance of any one of claims 23 to 30 to screen for binding partners to the substance.
57. A method of screening for substances which mimic the activity of BRCA2 polypeptide or a portion thereof, the method comprising contacting the test substances with a BRCA2 specific binding partner and, and determining whether the test substances bind to the specific binding partner.
58. The method of claim 57 further comprising the testing the substances binding to the specific binding partner for activity as cancer therapeutics.
59. A method of screening for substances which affect or modulate the activity of the BRCA2 polypeptide of any one of claims 23 to 30, the method comprising contacting one or more test substances with the BRCA2 polypeptide in a reaction medium, testing the activity of the treated BRCA2 polypeptide and comparing that activity with the activity of the BRCA2 polypeptide in comparable reaction medium untreated with the test substance

or substances.

60. A kit for detecting mutations in the BRCA2 gene associated with a susceptibility to cancer, the kit comprising one or more nucleic acid probes capable of specifically binding a mutated BRCA2 nucleic acid sequence.
61. A kit for detecting mutations in the BRCA2 gene associated with a susceptibility to cancer, the kit comprising one or more antibodies capable of specifically binding a mutated BRCA2 nucleic acid sequence.
62. A kit comprising at least one oligonucleotide primer having a sequence corresponding to or complementary to a portion of the nucleic acid sequence set out in any one of figures 1, 2, 4 or 7 for use in amplifying a BRCA2 nucleic acid sequence or an allele thereof.
63. The kit of claim 62 wherein the primers are for detecting mutations in the BRCA2 gene by single stranded conformation polymorphism (SSCP) and/or by a protein truncation (PTT) test.
64. A kit for determining the presence of one or more mutations in a sample of nucleic acid from an individual, the kit comprising:
- (a) a solid support having immobilised thereon one or more allele specific nucleic acid probes having sequences corresponding to portions of the sequence set out of figures 1, 2, 4 or 7 or a complementary sequence thereof and/or one or more antibodies capable of specifically binding a mutated BRCA2 nucleic acid sequence; and,
  - (b) a label for marking the presence of sample nucleic acid hybridised to the probe(s) or antibodies, or to probes or antibodies not hybridised to sample nucleic acid.
65. The kit of claim 64 wherein the label is adapted for binding to the nucleic acid sample prior to contacting the sample with the support.
66. The kit of claim 64 wherein the label is associated with a developing agent, the developing agent being capable of binding to sample nucleic acid hybridised to the probe(s) or antibodies, or to probes or antibodies not hybridised to the sample nucleic acid.
67. A chimeric animal having an normal BRCA2 allele.
68. A chimeric animal having a BRCA2 allele having one or more of the mutations set out table 1.

1/52

## Fig. 1.

Sequence 1. The BRCA2 cDNA

The sequence is shown 5' to 3'

```
NCNAAAGTTTAAAGGAGTTGTTTAGGAGGAAATTTGATTTAATTCAGAAA
NTGAGCATAGTNTTCCANTATTCACCTACGTTTTAGACCAAAATGTATCA
AAAATACTTCCTTCGTGNTGATAAGAGAAACCCAGAGCACTGTGTAAACT
CCAGAAATGGAAAAAACCTGCAGTAAAGAATTTAAATTATCAAATAACTT
AAATGTTGAAGGTGGTTNTTCAGAAAATAATCACTCTATTAAAGTTTCTC
CATATCTCTCTCAATTTCAACAAGACAAAACAACAGTTGGTATTAGGAACC
AARGTNTCACTTGGTTGAGAACATTCATGTTTTGGGAAAAGAACAGGCTTC
ACCTAAAAACGTAAAAATGGAAATTTGGTAAACTGAAACTTTTTCTGATG
TTCCTGTGNAAACAAATATAGAAGTTTGTCTACTTACTCCAAAGATTCA
GAAACTACTTTGAAACAGAAGCAGTAGAAATTGCTAAAGCTTTTATGGA
AGATGATGAAGTACAGATTNTAACTGCCAAGTCATGCCACACATNTC
TTTTTACATGTCCCGAAAAATGAGGAAATGGTTCTGTCAAATTCAGAATT
GGAAAAGAAGAGGAGAGCCCTTATCTTAGTGGGAGAACCCTCAATCAA
AAGAACTTATTAAATGAATTTGACAGGATAATAGAAAATCAAGAAAAAT
CCTTAAAGGCTTCAAAAAGCACTCCAGATGGCACAATAAAAGATTGAAGA
TTGTTTGTGCATCATGTTTCTTTAGAGCCGATTACCTGTGTACCCTTTTCG
CACACTAAGGAACCTCAAGAGATACAGAATCCAAATTTTACCGCACCTG
GTCAAGAATTTCTGTCTAAATCTCATTGTGTATGAACATCTGACTTTGGAA
AAATCTTCAAGCAATTTAGCAGTTTCAGGACATCCATTTTATCAAGTTTC
TGCTACAAGAAATGAAAAAATGAGACACTTGATTACTACAGGCAGACCGA
CCAAAGTCTTTGTTCCACCTTTTAAAACTAAATCACATTTTCACAGAGTT
GAACAGTGTGTTAGGAATATTAACTTGGAGGAAAACAGACAAAAGCAAAA
CATTGATGGACATGGCTCTGATGATAGTAAAAATAAGATTAATGACAATG
AGATTCATCAGTTTAAACAAAAACAACCTCCAATCAAGCAGCAGCTGTAAC
TTCACAAAGTGTGAAGAAGAACCTTTAGATTTAATTACAAGTCTTCAGAA
TGCCAGAGATATACAGGATATGCGAATTAAGAAGAAACAAAGGCAACCGG
TCTTTCCACAGCCAGGCAGTCTGTATCTTGCAAAAACATCCACTCTGCCT
CGAATCTCTCTGAAAGCAGCAGTAGGAGGCCAAGTTCCCTCTGCGTGTTC
TCATAAACAGCTGTATACGTATGGCGTTTCTAAACATTGCATAAAAAATTA
ACAGCAAAAATGCAGAGTCTTTTCAGTTTCACACTGAAGATTATTTTGGT
AAGGAAAGTTTATGGACTGGAAAAGGAATACAGTTGGCTGATGGTGGATG
GCTCATACCCTCCAATGATGGAAAGGCTGGAAAAGAAGAATTTTATAGGG
CTCTGTGTGACGTAAAGGCCACATAGTGGATAAACCATGTGCCACTTGAG
GAATTGAGAAAAAGTCAGTGTGTTTAGAACATAAGGAGAGGGGCAGAGAA
ATGGATCTGGTACAGGAGAATCAATGAGACCGGGTAAGAAACAGAAAAGG
GGCTGCACCAATGATTGACACGCTCTGCAAACTTCTTTTGTGGCTCAA
GTTGTGGCTCAAGAGGTGAGAAGGTAAGGCCAGACTTATTTGTTATGATT
TGCCCTTTAAATTGAAGCCTTAAGATTGGAATTCGATATCAAGCTTATCG
ATACCGTCGACCTCGAG
```



2/52

## Fig.2(a).

Genomic organisation of the BRCA2 gene

exon29f

genomic cgtaaaaatggaaattggtaaaaactgaaactttttctgatgttcctgtga

genomic anncaaanatagaagtttgttctacttacncccaagattcagaaaactac

genomic tttgnaacagaancagnagaaattgctaaagcttttatggaagatgatga

genomic actgacagattctaantctccaantcatgccacacattctctttttacat

cDNA GTCCCGAAAATGAGGAAATGGTTTAGTCAAATTCAAGAATTGGAAAAAGA

genomic GTCCCGAAAATGAGGAAATGGTTTGTGTCNAANTCAAGAATTGGAAAAAGA

cDNA AGAGGAGAGCCCCTTATCTTA-GTGG

genomic AGAGGAGAGCCCCTTATCTTANCNGGgtaagtnttcattttttaccttttcgcc

genomic rttgccaatcactattttttaaagtgtttattcagtagacttggtatgcta

genomic acaattaagagtgttataaactatgtcttttcagccatttttgtgtagtc

genomic agtttgggggagtatggtttgatatacagatacacagattcagtatcgt

genomic atacagatttgatatcttggtatacagattcgatatctctgaatctgtat

3/52

## Fig.2 (b).

exon29e

cDNA GAGAACCCTCAATCAAAAGAACTTATTAAATGAATTTGACAGGATAAT

cDNA A.GAAAATCAAGAAAAATCCTTAAAGGCTTCAAAAAGCACTCCAGATG  
 genomic AAATcAaGAAAA-TCCTTAAAGGNTTCAAAAAGCACTCCAGATG

genomic gtaaaattagctttttatattatctgttctccctctataggtatgggtat

genomic ataatattctgacctcaggtgatccacctgcctctcaaagtgcgtgggatt

genomic acagacatgagccactgtgcctaatacaagggacctctttataactctttaa

genomic aattactgaggacctaaaagagcatttggttatgtggaatatatctattg

## Fig.2 (c).

exon29d

genomic tgtactgtgagttatttggtgcatagtcattatcaatttgtgaatcaatt

genomic tattttcatagttaacatttattgagcatccgttacattcactgaaaatt

genomic gttaaagcctataaattgtctcaaattttttgtgtatttacagtaacatgga

genomic tattctcttagattttaactaatatgtaatatataaaataattgtttcctag

cDNA GCACAATAAAAAGATCGAAGATTGTTTATGCATCATGTTTCTTTAGAGCCG  
 genomic GCACAATAAAAAGATCGAAGATTGTAGGGCATCAGTT AGCCG

cDNA ATTACCTGTGTACCCTTTTCG

genomic ATTACCTGTGTACCCTTTTCGgtaagacatgtttaaatttttctaaattct

genomic aatacagtatgagaaaagtctcggtttttataaatgaacatttctaaaaat

genomic aatgacactaacgttaagaagttaacacttcccggttttataaaatttata

genomic aaatactttggtagtattttatagtgctgttcatatcattattttatttt

genomic ttaattttatgacagctttgtaaagtagacagattttattctaattttat

4/52

## Fig.2(d).

exon bc

genomic acacactgagaggtccacacttgacagatttatntattatgngttatgt

genomic gaggtagattgtaagtcaaaggctagccttgaaaaatgtgatattgtttt

genomic ggaatggcaaccatggngaanaaaaaacagtnaccagaatngtatcacca

genomic tgtagcaaatgagggtctgcaacaaaggcatattcataaatattnanang

cDNA CACAACTAA  
genomic tgtagtagtcaatnaacttataaattttntccccantgcagCACNANTAA

cDNA GGAACGTCAAGAGATACAGAATCCAAATTTTACCGCACCTGGTCAAGAAT  
genomic GGGACGTCAAGAGATACAGAATCCANATTTTACCGCACCTGGTCANGGAT

cDNA TTCTGTCTAAATCTCATTGTGTATGAACATCTGACTTTGGAAAAATCTTCA  
genomic TTCTGTNTAAATATCATTGTGTATGAACATCTGACTTTGGAAAAATCTTCA

cDNA AGCAATTTAGCAGTTTCAGGACATCCATTTTATCAAGTTTCTGCTACAAG  
genomic AGCAATTTNGCAGTTTCAGGACATCCATTTTATCAAGTTTCTGCTACAAG

cDNA AAATGAAAAAATGAGACACTTGATTACTACAGGCAGACCAACCAAAGTCT  
genomic AAATGAAAAAATGAGACACTTGATTACTACAGGCAGACCAACCAAAGTCT

cDNA TTGTTCCACCTTTTAAAACTAAATCACATTTTC-ACAGAGTTGAACAGTGT  
genomic TTGTTCCACCTTTTAAAACTAAATCACATTTTCCACAGAGTTGAACAGTGT

cDNA GTTAGGAATATTAACCTTGGAGGAAAACAGACAAAAGCAAAACATTGATGG  
genomic ggangaacatATTNAAGCAAAACATTGATGG

cDNA ACATGGCTCTGATGATAGTAAAAATAAGATTAATGACAATGAGATTCATC  
genomic ACNTGGCTCTGATGATAGTAAAAATAAGATTAATGACAATGAGATTCATC

cDNA AGTTTAAACAGNAACAACTCCAATCAAGCAGCAGCTGTAACCTTTCACAAAG  
genomic AGTTTAAACAAAAACAACTCCAATCAAGCAGCAGCTGTAACCTTTCACAAAG

cDNA TGTGAAGAAGAACCTTTAG  
genomic TGTGAANAAAAACCTTTAGgtattgtatgacaatttgtgtgatgaatttt

genomic tgcctttcagttanatattccggttgtaaataatgtcctgatgggtttnc

5/52

## Fig.2 (e).

genomic cccctttggtggtggttaattttaagccctttttaatgttttagattttc

genomic taaatccaaagattaggtttaaattattctaagtgttctttcaaanataa

exon a

cDNA ATTTAATTACAAGTCTTCAGAATGCCAGAGA  
cDNA TATACAGGATATGCGAATTAAGAAGAAACAAAGGCAACGCGTCTTCCAC  
cDNA AGCCAGGCAGTCTGTATCTTGCAAAAACATCCACTCTGCCTCGAATCTCT  
cDNA CTGAAAGCAGCAGTAGGAGGCCAAGTTCCTCTGCGTGTTCATATAACA  
cDNA GCTGTATACGTATGGCGTTTCTAAACATTGCATAAAAATTAACAGCAAAA  
cDNA ATGCAGAGTCTTTTCAGTTTCACACTGAAGATTATTTTGGTAAGGAAAGT  
cDNA TTATGGACTGGAAAAGGAATACAGTTGGCTGATGGTGGATGGCTCATACC  
cDNA CTCCAATGATGGAAAGGCTGGAAAAGAAGAATTTTATAGGGCTCTGTGTG  
cDNA ACGTAAAGGCCACATAGTGGATAAACCATGTGCCACTTGAGGAATTGAGA  
cDNA AAAAGTCAGTGTGTTTAGAACATAAGGAGAGGGGCAGAGAAATGGATCTG  
cDNA GTACAGGAGAATCAATGAGACCGGGTAAGAAACAGAAAAGGGCTGCACC  
cDNA AAATGATTGACACGCTCTGCAAACCTCTTTTGTGGCTCAAGTTGTGGCT  
cDNA CAAGAGGTGAGAAGGTAAGGCCAGACTTATTTGTTATGATTTGCCCTTTA  
cDNA AATTGAAGCCTTAAGATTGGAATTCGATATCAAGCTTATCGATACCGTCG  
cDNA ACCTCGAG

6/52

## Fig.3.

## BRCA2 protein sequence

Plaque 14 was translated in all 3 possible frames. The following protein sequence is the most likely of the 3 frames.

KVGSCLGGNLIFRXAXSXIHLRFRPKCIKNTSFVXIRETQSTVTPEMEKT  
CSKEFKLSNNLNVEGGXSENNHSIKVSPYLSQFQQDKQQLVLGTVSLVE  
NIHVLGKEQASPKNVKMEIGKTETFSQVVPVXTNIEVCSTYSKDSENYFET  
EAVEIAKAFMEDDELTDXKLPSHATHXLFTCPENEEMVLNSRIGKRRGE  
PLILVGEPSTIKRNLLNEFDRIIENQEKSLKASKSTPDGTIKDRLFVHHVS  
LEPITCVPFRTTKERQEIQNPNFTAPGQEFSLKSHLYEHLTLEKSSSNLA  
VSGHPFYQVSATRNEKMRHLITTGRPTKVFPVPPFKTKSHFHRVEQCVRNI  
NLEENRQKQONIDGHGSDDSKNKINDNEIHQFNKNNSNQAAAVTFTKCEE  
PLDLITSLQNARDIQDMRIKKKQORQVFPQPGSLYLAKTSTLPRI SLKAA  
VGGQVPSACSHKQLYTYGVSKHC IKINSKNAESFQFHTEDYFGKESLWTG  
KGIQLADGGW LIPSN DGKAGKEEFYRALCDVKAT

7/52

## Fig.4(a).

BRCA2 cDNA

CCACATTGGAAAGTCAATGCCAAATGTCCTAGAAGATGAAGTATATGAAACAGTTGTAGA  
TACCTCTGAAGAAGATAGTTTTTCATTATGTTTTTCTAAATGTAGAACAAAAATCTACA  
AAAAGTAAGAACTAGCAAGACTAGGAAAAAAATTTTCCATGAAGCAAACGCTGATGAATG  
TGAAAAATCTAAAAACCAAGTGAAAGAAAAATACTCATTTGTATCTGAAGTGGAACCAAA  
TGATACTGATCCATTAGATTCAAATGTAGCAAATCAGAAGCCCTTTGAGAGTGGAAGTGA  
CAAAATCTCCAAGGAAGTTGTACCGTCTTTGGCCTCTGAATGGTCTCAACTAACCCCTTTC  
AGGTCTAAATGGAGCCCAGATGGAGAAAAATACCCCTATTGCATATTTCTTCATGTGACCA  
AAATATTTTCAGAAAAAGACCTATTAGACACAGAGAACAAAAGAAAGAAAGATTTTCTTAC  
TTCAGAGAAATCTTTGCCACGTATTTCTAGCCTACCAAAATCAGAGAAGCCATTAAATGA  
GGAAACAGTGGTAAATAAGAGAGATGAAGAGCAGCATCTTGAATCTCATACAGACTGCAT  
TCTTGCAGTAAAGCAGGCAATATCTGGAACCTCTCCAGTGGCTTCTTCATTTTCAGGGTAT  
CAAAAAGTCTATATTCAGAATAAGAGAATCACCTAAAGAGACTTTCAATGCAAGTTTTTC  
AGGTCTATGACTGATCCAACTTTAAAAAGAAACTGAAGCCTCTGAAAGTGGACTGGA  
AATACATACTGTTTGTCTCACAGAAGGAGGACTCCTTATGTCCAAATTTAATTGATAATGG  
AAGCTGGCCAGCCACCACCACACAGAATTCTGTAGCTTTGAAGAATGCAGGTTTAATATC  
CACTTTGAAAAAGAAAAACAAATAAGTTTATTTATGCTATACATGATGAAACATCTTATAA  
AGGAAAAAAAATACCGAAAGACCAAAAATCAGAACTAATTAAGTCTTCAGCCCAGTTTGA  
AGCAATGCTTTTGAAGCACCACCTTACATTTGCAAATGCTGATTGAGGTTTATTGCATTC  
TTCTGTGAAAAGAAGCTGTTTACAGAATGATTCTGAAGAACCAACTTTGTCTTAAGTAG  
CTCTTTTGGGACAATTCTGAGGAAATGTTCTAGAAATGAAACATGTTCTAATAATACAGT  
AATCTCTCAGGATCTTGATTATAAAGAAGCAAAATGTAATAAGGAAAAACTACAGTTATT  
TATTACCCCGAAGCTGATTCTCTGTGCATGCCTGCAGGAAGGACAGTGTGAAAATGATCC  
AAAAAGCAAAAAGTTTCAGATATAAAGAAGAGGTCTTGGCTGCAGCATGTCACCCAGT  
ACAACATTCAAAGTGGAATACAGTGATACTGACTTTCAATCCCAGAAAAGTCTTTTATA  
TGATCATGAAAATGCCAGCACTCTTATTTTAACCTCCTACTTCCAAGGATGTTCTGTCAAA  
CCTAGTCATGATTTCTAGAGGCAAAGAATCATACAAAATGTCAGACAAGCTCAAAGGTAA  
CAATTATGAATCTGATGTTGAATTAACCAAAAATATTTCCCATGGAAAAGAATCAAGATGT  
ATGTGCTTTAATGAAAATTATAAAAACGTTGAGCTGTTGCCACCTGAAAAATACATGAG

8/52  
Fig.4(b).

AGTAGCATCACCTTCAAGAAAGGTACAATTCAACCAAAACACAAATCTAAGAGTAATCCA  
AAAAAATCAAGAAGAACTACTTCAATTTCAAAAATAACTGTCAATCCAGACTCTGAAGA  
ACTTTTCTCAGACAATGAGAATAATTTTGTCTTCCAAGTAGCTAATGAAAGGAATAATCT  
TGCTTTAGGAAATACTAAGGAACCTTCATGAAACAGACTTGACTTGTGTAAACGAACCCAT  
TTTCAAGAACTCTACCATGGTTTTATATGGAGACACAGGTGATAAACAAGCAACCCAAGT  
GTCAATTAAAAAAGATTTGGTTTATGTTCTTGCAGAGGAGAACAAAAATAGTGTAAGCA  
GCATATAAAAAATGACTCTAGGTCAAGATTTAAAAATCGGACATCTCCTTGAATATAGATAA  
AATACCAGAAAAAATAATGATTACATGAACAAATGGGCAGGACTCTTAGGTCCAATTTTC  
AAATCACAGTTTTGGAGGTAGCTTCAGAACAGCTTCAAATAAGGAAATCAAGCTCTCTGA  
ACATAACATTAAGAAGAGCAAAATGTTCTTCAAAGATATTGAAGAACAATATCCTACTAG  
TTTAGCTTGTGTTGAAATTGTAAATACCTTGGCATTAGATAATCAAAAGAACTGAGCAA  
GCCTCAGTCAATTAATACTGTATCTGCACATTTACAGAGTAGTGTAGTTGTTTCTGATTG  
TAAAAATAGTCATATAACCCCTCAGATGTTATTTTCCAAGCAGGATTTTAATTCAAACCA  
TAATTTAACACCTAGCCAAAAGGAGCAAATTACAGAACTTTCTACTATATTAGAAGATTC  
AGGAAGTCAGTTTGAATTTACTCAGTTTAGAAAACCAAGCTACATATTGCAGAAGAGTAC  
ATTTGAAGTGCCTGAAAACCAGATGACTATCTTAAAGACCACTTCTGAGGAATGCAGAGA  
TGCTGATCTTCATGTCATAATGAATGCCCCATCGATTGGTCAGGTAGACAGCAGCAAGCA  
ATTTGAAGGTACAGTTGAAATTAAACGGAAGTTTGCTGGCCTGTTGAAAAATGACTGTAA  
CAAAAGTGCTTCTGGTTATTTAACAGATGAAAATGAAGTGGGGTTTAGGGGCTTTTATTC  
TGCTCATGGCACAAAACCTGAATGTTTCTACTGAAGCTCTGCAAAAAGCTGTGAAACTGTT  
TAGTGATATTGAGAATATTAGTGAGGAACTTCTGCAGAGGTACATCCAATAAGTTTATC  
TTCAAGTAAATGTCATGATTCTGTCGTTTCAATGTTTAAGATAGAAAATCATAATGATAA  
AACTGTAAGTGAAAAAATAATAAATGCCAACTGATATTACAAAATAATATTGAAATGAC  
TACTGGCACTTTTGTGTTGAAGAAATTACTGAAAATTACAAGAGAAATACTGAAAATGAAGA  
TAACAAATATACTGCTGCCAGTAGAAATTCTCATAACTTAGAATTTGATGGCAGTGATTC  
AAGTAAAAATGATACTGTTTGTATTCATAAAGATGAAACGGACTTGCTATTTACTGATCA  
GCACAACATATGTCTTAAATTATCTGGCCAGTTTATGAAGGAGGGAAACACTCAGATTAA  
AGAAGATTTGTCAGATTTAACTTTTTTGAAGTTGCGAAAGCTCAAGAAGCATGTCATGG  
TAATACTTCAAATAAAGAACAGTTAACTGCTACTAAAACGGAGCAAAATATAAAAGATTT

9/52

## Fig.4(c).

TGAGACTTCTGATACATTTTTTCAGACTGCAAGTGGGAAAAATATTAGTGTGCGCCAAAGA  
GTCATTTAATAAAATTGTAAATTTCTTTGATCAGAAACCAGAAGAATTGCATAACTTTTC  
CTTAAATTCTGAATTACATTCTGACATAAGAAAGAACAAAATGGACATTCTAAGTTATGA  
GGAAACAGACATAGTTAAACACAAAATACTGAAAGAAAGTGTCCCAGTTGGTACTGGAAA  
TCAACTAGTGACCTTCCAGGGACAACCCGAACGTGATGAAAAGATCAAAGAACCTACTCT  
GTTGGGTTTTTCATACAGCTAGCGGGAAAAAAGTTAAATTGCAAAGGAATCTTTGGACAA  
AGTGAAAACCTTTTTGATGAAAGAGCAAGGTACTAGTGAAATCACCAGTTTTAGCCATC  
AATGGGCAAAGACCCTAAAGTACAGAGAGGCCTGTAAAGACCTTGAATTAGCATGTGAGA  
CCATTGAGATCACAGCTGCCCCAAAGTGTAAGAAATGCAGAATTCTCTCAATAATGATA  
AAAACCTTGTTTCTATTGAGACTGTGGTGCCACCTAAGCTCTTAAGTGATAATTTATGTA  
GACAACTGAAATCTCAAAACATCAAAAAGTATCTTTTGAAGTTAAAGTACATGAAA  
ATGTAGAAAAAGAAACAGCAAAAAGTCCTGCAACTTGTTACACAAATCAGTCCCCTTATT  
CAGTCATTGAAAATTCAGCCTTAGCTTTTTACACAAGTTGTAGTAGAAAACTTCTGTGA  
GTCAGACTTCATTACTTGAAGCAAAAAAATGGCTTAGAGAAGGAATATTTGATGGTCAAC  
CAGAAAGAATAAATACTGCAGATTATGTAGGAAATTATTTGTATGAAAATAATTCAAACA  
GTACTATAGCTGAAAATGACAAAATCATCTCTCCGAAAAACAAGATACTTATTTAAGTA  
ACAGTAGCATGTCTAACAGCTATTCCCTACCATTCTGATGAGGTATATAATGATTCAGGAT  
ATCTCTCAAAAAATAAACTTGATTCTGGTATTGAGCCAGTATTGAAGAATGTTGAAGATC  
AAAAAACACTAGTTTTTCCAAAGTAATATCCAATGTAAAAGATGCAAATGCATACCCAC  
AAACTGTAAATGAAGATATTTGCGTTGAGGAACTTGTGACTAGCTCTTCACCCTGCAAAA  
ATAAAAATGCAGCCATTAAATTGTCCATATCTAATAGTAATAATTTGAGGTAGGGCCAC  
CTGCATTTAGGATAGCCAGTGGTAAATCGTTTGTGTTTCACATGAAACAATTAAAAAAG  
TGAAAGACATATTTACAGACAGTTTCAGTAAAGTAATTAAGGAAAACAACGAGAATAAAT  
CAAAAAATTTGCCAAACGAAAATTATGGCAGGTTGTTACGAGGCATTGGATGATTCAGAGG  
ATATTCTTCATAACTCTCTAGATAATGATGAATGTAGCACGCATTACATAAGGTTTTTG  
CTGACATTCAGAGTGAAGAAATTTTACAACATAACCAAAATATGTCTGGATTGGAGAAAG  
TTTCTAAAATATCACCTTGATGTTAGTTTGGAACTTCAGATATATGTAAATGTAGTA  
TAGGGAAGCTTCATAAGTCAGTCTCATCTGCAAATACTTGTGGGATTTTTAGCACAGCAA  
GTGGAAAATCTGTCCAGGTATCAGATGCTTCATTACAAAACGCAAGACAAGTGTTCCTG



10/52

## Fig.4(d).

AAATAGAAGATAGTACCAAGCAAGTCTTTTCCAAAGTATTGTTTAAAAGTAACGAACATT  
CAGACCAGCTCACAAGAGAAGAAAATACTGCTATACGTACTCCAGAACATTTAATATCCC  
AAAAAGGCTTTTCATATAATGTGGTAAATTCATCTGCTTTCTCTGGATTTAGTACAGCAA  
GTGGAAAGCAAGTTTCCATTTTAGAAAAGTTCCTTACACAAAGTTAAGGGAGTGTTAGAGG  
AATTTGATTTAATCAGAACTGAGCATAGTCTTCACTATTACCTACGTCTAGACAAAATG  
TATCAAAAATACTTCCTCGTGTGATAAGAGAAAACCCAGAGCACTGTGTAAACTCAGAAA  
TGGAAAAAACCTGCAGTAAAGAATTTAAATTATCAAATAACTTAAATGTTGAAGGTGGTT  
CTTCAGAAAATAATCACTCTATTAAAGTTTCTCCATATCTCTCTCAATTTCAACAAGACA  
AACACAGTTGGTATTAGGAACCAAAGTCTCACTTGTTGAGAACATTCATGTTTTGGGAA  
AAGAACAGGCTTCACCTAAAAACGTAAAAATGGAAATTGGTAAACTGAACTTTTTCTG  
ATGTTCTGTGAAAACAAATATAGAAGTTTGTCTACTTACTCCAAAGATTCAGAAAAT  
ACTTTGAAACAGAAGCAGTAGAAATTGCTAAAGCTTTTATGGAAGATGATGAACTGACAG  
ATTCTAAACTGCCAAGTCATGCCACACATTCTCTTTTACATGTCCCGAAAATGAGGAAA  
TGGTTTTGTCAAATTCAAGAATTGGAAAAAGAAGAGGAGAGCCCTTATCTTAGTGGGAG  
AACCCTCAATCAAAAGAACTTATTAAATGAATTTGACAGGATAATAGAAAATCAAGAAA  
AATCCTTAAAGGCTTCAAAAAGCACTCCAGATGGCACAATAAAAGATCGAAGATTGTTTG  
TGCATCATGTTTCTTTAGAGCCGATTACCTGTGTACCCCTTTCGCACAACCTAAGGAACGTC  
AAGAGATACAGAATCCAAATTTTACCGCACCTGGTCAAGAATTTCTGTCTAAATCTCATT  
TGTATGAACATCTGACTTTGGAAAAATCTTCAAGCAATTTAGCAGTTTCAGGACATCCAT  
TTTATCAAGTTTCTGGTAACAAGAATGGAAAAATGAGAAAATTGATTACTACAGGCAGAC  
CAACCAAAGTCTTTGTTCCACCTTTTAAACTAAATCACATTTTCACAGAGTTGAACAGT  
GTGTTAGGAATATTAACCTGGAGGGAAACAGACAAAACCAAACATTGATGGACATGGCT  
CTGATGATAGTAAAAATAAGATTAATGACAATGAGATTCATCAGTTTAACAAAAACAAC  
CCAATCAAGCAGCAGCTGTAACTTTCACAAAGTGTGAAGAAGAACCTTTAGATTTAATTA  
CAAGTCTTCAGAAATGCCAGAGATATACAGGATATGCGAATTAAGAAGAAACAAAGGCAAC  
GCGTCTTTCCACAGCCAGGCAGTCTGTATCTTGCAAAAACATCCACTCTGCCTCGAATCT  
CTCTGAAAGCAGCAGTAGGAGGCCAAGTTCCCTCTGCGTGTCTCATAAACAGCTGTATA  
CGTATGGCGTTTCTAAACATTGCATAAAAATTAACAGCAAAAATGCAGAGTCTTTTCAGT  
TTCACACTGAAGATTATTTTGGTAAGGAAAGTTTATGGACTGGAAAAGGAATACAGTTGG

SUBSTITUTE SHEET (RULE 26)

11/52

Fig.4(e).

CTGATGGTGGATGGCTCATACCCTCCAATGATGGAAAGGCTGGAAAAGAAGAATTTTATA  
GGGCTCTGTGTGACGTAAAGGCCACATAGTGGATAAACCATGTGCCACTTGAGGAATTGA  
GAAAAAGTCAGTGTGTTTAGAACATAAGGAGAGGGGCAGAGAAATGGATCTGGTACAGGA  
GAATCAATGAGACCGGGTAAGAAACAGAAAAGGGGCTGCACCAAATGATTGACACGCTCT  
GCAAACCTCTTTTGTGGCTCAAGTTGTGGCTCAAGAGGTGAGAAGGTAAGGCCAGACTT  
ATTTGTTATGATTTGCCCTTTAAATTGAAGCCTTAAGATT

12/52

Fig.5(a).

HisIleGlyLysSerMetProAsnValLeuGluAspGluValTyrGluThrValValAsp 20  
 ThrSerGluGluAspSerPheSerLeuCysPheSerLysCysArgThrLysAsnLeuGln 40  
 LysValArgThrSerLysThrArgLysLysIlePheHisGluAlaAsnAlaAspGluCys 60  
 GluLysSerLysAsnGlnValLysGluLysTyrSerPheValSerGluValGluProAsn 80  
 AspThrAspProLeuAspSerAsnValAlaAsnGlnLysProPheGluSerGlySerAsp 100  
 LysIleSerLysGluValValProSerLeuAlaCysGluTrpSerGlnLeuThrLeuSer 120  
 GlyLeuAsnGlyAlaGlnMetGluLysIleProLeuLeuHisIleSerSerCysAspGln 140  
 AsnIleSerGluLysAspLeuLeuAspThrGluAsnLysArgLysLysAspPheLeuThr 160  
 SerGluAsnSerLeuProArgIleSerSerLeuProLysSerGluLysProLeuAsnGlu 180  
 GluThrValValAsnLysArgAspGluGluGlnHisLeuGluSerHisThrAspCysIle 200  
 LeuAlaValLysGlnAlaIleSerGlyThrSerProValAlaSerSerPheGlnGlyIle 220  
 LysLysSerIlePheArgIleArgGluSerProLysGluThrPheAsnAlaSerPheSer 240  
 GlyHisMetThrAspProAsnPheLysLysGluThrGluAlaSerGluSerGlyLeuGlu 260  
 IleHisThrValCysSerGlnLysGluAspSerLeuCysProAsnLeuIleAspAsnGly 280  
 SerTrpProAlaThrThrThrGlnAsnSerValAlaLeuLysAsnAlaGlyLeuIleSer 300  
 ThrLeuLysLysLysThrAsnLysPheIleTyrAlaIleHisAspGluThrSerTyrLys 320  
 GlyLysLysIleProLysAspGlnLysSerGluLeuIleAsnCysSerAlaGlnPheGlu 340  
 AlaAsnAlaPheGluAlaProLeuThrPheAlaAsnAlaAspSerGlyLeuLeuHisSer 360  
 SerValLysArgSerCysSerGlnAsnAspSerGluGluProThrLeuSerLeuThrSer 380  
 SerPheGlyThrIleLeuArgLysCysSerArgAsnGluThrCysSerAsnAsnThrVal 400  
 IleSerGlnAspLeuAspTyrLysGluAlaLysCysAsnLysGluLysLeuGlnLeuPhe 420  
 IleThrProGluAlaAspSerLeuSerCysLeuGlnGluGlyGlnCysGluAsnAspPro 440  
 LysSerLysLysValSerAspIleLysGluGluValLeuAlaAlaAlaCysHisProVal 460  
 GlnHisSerLysValGluTyrSerAspThrAspPheGlnSerGlnLysSerLeuLeuTyr 480  
 AspHisGluAsnAlaSerThrLeuIleLeuThrProThrSerLysAspValLeuSerAsn 500  
 LeuValMetIleSerArgGlyLysGluSerTyrLysMetSerAspLysLeuLysGlyAsn 520  
 AsnTyrGluSerAspValGluLeuThrLysAsnIleProMetGluLysAsnGlnAspVal 540  
 CysAlaLeuAsnGluAsnTyrLysAsnValGluLeuLeuProProGluLysTyrMetArg 560  
 ValAlaSerProSerArgLysValGlnPheAsnGlnAsnThrAsnLeuArgValIleGln 580  
 LysAsnGlnGluGluThrThrSerIleSerLysIleThrValAsnProAspSerGluGlu 600  
 LeuPheSerAspAsnGluAsnAsnPheValPheGlnValAlaAsnGluArgAsnAsnLeu 620  
 AlaLeuGlyAsnThrLysGluLeuHisGluThrAspLeuThrCysValAsnGluProIle 640  
 PheLysAsnSerThrMetValLeuTyrGlyAspThrGlyAspLysGlnAlaThrGlnVal 660  
 SerIleLysLysAspLeuValTyrValLeuAlaGluGluAsnLysAsnSerValLysGln 680  
 HisIleLysMetThrLeuGlyGlnAspLeuLysSerAspIleSerLeuAsnIleAspLys 700  
 IleProGluLysAsnAsnAspTyrMetAsnLysTrpAlaGlyLeuLeuGlyProIleSer 720  
 AsnHisSerPheGlyGlySerPheArgThrAlaSerAsnLysGluIleLysLeuSerGlu 740  
 HisAsnIleLysLysSerLysMetPhePheLysAspIleGluGluGlnTyrProThrSer 760

13/52

Fig.5(b).

LeuAlaCysValGluIleValAsnThrLeuAlaLeuAspAsnGlnLysLysLeuSerLys 780  
 ProGlnSerIleAsnThrValSerAlaHisLeuGlnSerSerValValValSerAspCys 800  
 LysAsnSerHisIleThrProGlnMetLeuPheSerLysGlnAspPheAsnSerAsnHis 820  
 AsnLeuThrProSerGlnLysGluGlnIleThrGluLeuSerThrIleLeuGluAspSer 840  
 GlySerGlnPheGluPheThrGlnPheArgLysProSerTyrIleLeuGlnLysSerThr 860  
 PheGluValProGluAsnGlnMetThrIleLeuLysThrThrSerGluGluCysArgAsp 880  
 AlaAspLeuHisValIleMetAsnAlaProSerIleGlyGlnValAspSerSerLysGln 900  
 PheGluGlyThrValGluIleLysArgLysPheAlaGlyLeuLeuLysAsnAspCysAsn 920  
 LysSerAlaSerGlyTyrLeuThrAspGluAsnGluValGlyPheArgGlyPheTyrSer 940  
 AlaHisGlyThrLysLeuAsnValSerThrGluAlaLeuGlnLysAlaValLysLeuPhe 960  
 SerAspIleGluAsnIleSerGluGluThrSerAlaGluValHisProIleSerLeuSer 980  
 SerSerLysCysHisAspSerValValSerMetPheLysIleGluAsnHisAsnAspLys 1000  
 ThrValSerGluLysAsnAsnLysCysGlnLeuIleLeuGlnAsnAsnIleGluMetThr 1020  
 ThrGlyThrPheValGluGluIleThrGluAsnTyrLysArgAsnThrGluAsnGluAsp 1040  
 AsnLysTyrThrAlaAlaSerArgAsnSerHisAsnLeuGluPheAspGlySerAspSer 1060  
 SerLysAsnAspThrValCysIleHisLysAspGluThrAspLeuLeuPheThrAspGln 1080  
 HisAsnIleCysLeuLysLeuSerGlyGlnPheMetLysGluGlyAsnThrGlnIleLys 1100  
 GluAspLeuSerAspLeuThrPheLeuGluValAlaLysAlaGlnGluAlaCysHisGly 1120  
 AsnThrSerAsnLysGluGlnLeuThrAlaThrLysThrGluGlnAsnIleLysAspPhe 1140  
 GluThrSerAspThrPhePheGlnThrAlaSerGlyLysAsnIleSerValAlaLysGlu 1160  
 SerPheAsnLysIleValAsnPhePheAspGlnLysProGluGluLeuHisAsnPheSer 1180  
 LeuAsnSerGluLeuHisSerAspIleArgLysAsnLysMetAspIleLeuSerTyrGlu 1200  
 GluThrAspIleValLysHisLysIleLeuLysGluSerValProValGlyThrGlyAsn 1220  
 GlnLeuValThrPheGlnGlyGlnProGluArgAspGluLysIleLysGluProThrLeu 1240  
 LeuGlyPheHisThrAlaSerGlyLysLysValLysIleAlaLysGluSerLeuAspLys 1260  
 ValLysAsnLeuPheAspGluLysGluGlnGlyThrSerGluIleThrSerPheSerHis 1280  
 GlnTrpAlaLysThrLeuLysTyrArgGluAlaCysLysAspLeuGluLeuAlaCysGlu 1300  
 ThrIleGluIleThrAlaAlaProLysCysLysGluMetGlnAsnSerLeuAsnAsnAsp 1320  
 LysAsnLeuValSerIleGluThrValValProProLysLeuLeuSerAspAsnLeuCys 1340  
 ArgGlnThrGluAsnLeuLysThrSerLysSerIlePheLeuLysValLysValHisGlu 1360  
 AsnValGluLysGluThrAlaLysSerProAlaThrCysTyrThrAsnGlnSerProTyr 1380  
 SerValIleGluAsnSerAlaLeuAlaPheTyrThrSerCysSerArgLysThrSerVal 1400  
 SerGlnThrSerLeuLeuGluAlaLysLysTrpLeuArgGluGlyIlePheAspGlyGln 1420  
 ProGluArgIleAsnThrAlaAspTyrValGlyAsnTyrLeuTyrGluAsnAsnSerAsn 1440  
 SerThrIleAlaGluAsnAspLysAsnHisLeuSerGluLysGlnAspThrTyrLeuSer 1460  
 AsnSerSerMetSerAsnSerTyrSerTyrHisSerAspGluValTyrAsnAspSerGly 1480  
 TyrLeuSerLysAsnLysLeuAspSerGlyIleGluProValLeuLysAsnValGluAsp 1500  
 GlnLysAsnThrSerPheSerLysValIleSerAsnValLysAspAlaAsnAlaTyrPro 1520

14/52

## Fig.5(c).

GlnThrValAsnGluAspIleCysValGluGluLeuValThrSerSerSerProCysLys 1540  
 AsnLysAsnAlaAlaIleLysLeuSerIleSerAsnSerAsnAsnPheGluValGlyPro 1560  
 ProAlaPheArgIleAlaSerGlyLysIleValCysValSerHisGluThrIleLysLys 1580  
 ValLysAspIlePheThrAspSerPheSerLysValIleLysGluAsnAsnGluAsnLys 1600  
 SerLysIleCysGlnThrLysIleMetAlaGlyCysTyrGluAlaLeuAspAspSerGlu 1620  
 AspIleLeuHisAsnSerLeuAspAsnAspGluCysSerThrHisSerHisLysValPhe 1640  
 AlaAspIleGlnSerGluGluIleLeuGlnHisAsnGlnAsnMetSerGlyLeuGluLys 1660  
 ValSerLysIleSerProCysAspValSerLeuGluThrSerAspIleCysLysCysSer 1680  
 IleGlyLysLeuHisLysSerValSerSerAlaAsnThrCysGlyIlePheSerThrAla 1700  
 SerGlyLysSerValGlnValSerAspAlaSerLeuGlnAsnAlaArgGlnValPheSer 1720  
 GluIleGluAspSerThrLysGlnValPheSerLysValLeuPheLysSerAsnGluHis 1740  
 SerAspGlnLeuThrArgGluGluAsnThrAlaIleArgThrProGluHisLeuIleSer 1760  
 GlnLysGlyPheSerTyrAsnValValAsnSerSerAlaPheSerGlyPheSerThrAla 1780  
 SerGlyLysGlnValSerIleLeuGluSerSerLeuHisLysValLysGlyValLeuGlu 1800  
 GluPheAspLeuIleArgThrGluHisSerLeuHisTyrSerProThrSerArgGlnAsn 1820  
 ValSerLysIleLeuProArgValAspLysArgAsnProGluHisCysValAsnSerGlu 1840  
 MetGluLysThrCysSerLysGluPheLysLeuSerAsnAsnLeuAsnValGluGlyGly 1860  
 SerSerGluAsnAsnHisSerIleLysValSerProTyrLeuSerGlnPheGlnGlnAsp 1880  
 LysGlnGlnLeuValLeuGlyThrLysValSerLeuValGluAsnIleHisValLeuGly 1900  
 LysGluGlnAlaSerProLysAsnValLysMetGluIleGlyLysThrGluThrPheSer 1920  
 AspValProValLysThrAsnIleGluValCysSerThrTyrSerLysAspSerGluAsn 1940  
 TyrPheGluThrGluAlaValGluIleAlaLysAlaPheMetGluAspAspGluLeuThr 1960  
 AspSerLysLeuProSerHisAlaThrHisSerLeuPheThrCysProGluAsnGluGlu 1980  
 MetValLeuSerAsnSerArgIleGlyLysArgArgGlyGluProLeuIleLeuValGly 2000  
 GluProSerIleLysArgAsnLeuLeuAsnGluPheAspArgIleIleGluAsnGlnGlu 2020  
 LysSerLeuLysAlaSerLysSerThrProAspGlyThrIleLysAspArgArgLeuPhe 2040  
 ValHisHisValSerLeuGluProIleThrCysValProPheArgThrThrLysGluArg 2060  
 GlnGluIleGlnAsnProAsnPheThrAlaProGlyGlnGluPheLeuSerLysSerHis 2080  
 LeuTyrGluHisLeuThrLeuGluLysSerSerSerAsnLeuAlaValSerGlyHisPro 2100  
 PheTyrGlnValSerGlyAsnLysAsnGlyLysMetArgLysLeuIleThrThrGlyArg 2120  
 ProThrLysValPheValProProPheLysThrLysSerHisPheHisArgValGluGln 2140  
 CysValArgAsnIleAsnLeuGluGlyAsnArgGlnLysGlnAsnIleAspGlyHisGly 2160  
 SerAspAspSerLysAsnLysIleAsnAspAsnGluIleHisGlnPheAsnLysAsnAsn 2180  
 SerAsnGlnAlaAlaAlaValThrPheThrLysCysGluGluGluProLeuAspLeuIle 2200  
 ThrSerLeuGlnAsnAlaArgAspIleGlnAspMetArgIleLysLysLysGlnArgGln 2220  
 ArgValPheProGlnProGlySerLeuTyrLeuAlaLysThrSerThrLeuProArgIle 2240  
 SerLeuLysAlaAlaValGlyGlyGlnValProSerAlaCysSerHisLysGlnLeuTyr 2260  
 ThrTyrGlyValSerLysHisCysIleLysIleAsnSerLysAsnAlaGluSerPheGln 2280

15/52

## Fig.5(d).

PheHisThrGluAspTyrPheGlyLysGluSerLeuTrpThrGlyLysGlyIleGlnLeu 2300  
AlaAspGlyGlyTrpLeuIleProSerAsnAspGlyLysAlaGlyLysGluGluPheTyr 2320  
ArgAlaLeuCysAspValLysAlaThrStop 2329

16/52

**SUBSTITUTE SHEET (RULE 26)**

**SUBSTITUTE SHEET (RULE 26)**

Putative *BRCA2* promoter region. Potential binding sites for various transcription factors are underlined. CpG are emboldened.



18/52

Fig.7. (1/22)

the genomic sequence available from:  
ftp://sanger.ac.uk/pub/human/sequences/13q  
and ftp://genome.wustl.edu/pub/gsc1/brca2  
and is present in the Pacs 214k23 and 92m18

Exon	1	2	3	4	5	6	7	8	9
Size	188	106	249	109	50	41	115	50	112

Exon	10	11	12	13	14	15	16	17	18
Size	1116	4932	96	70	428	166	204	171	355

Exon	19	20	21	22	23	24	25	26	27
Size	156	145	122	199	164	139	245	147	606

NB there is 67bp of coding sequence in exon 2

Exon 1

.....aggggggncccccttcccaaaagnaag  
ggtggcccttggnaahtcctnaaaggntcaangcnaantaagaaagaaacacnaacaa  
ctcccaaattccccgctttaattcggtcaagaataactaacggttgggaatgccttgaacaa  
aggaagatttcccttcgcgaacactgagaaaaatacccgcaagcgcccnaccnaggcctga  
cttccgggttggtgcgtgtgctgcgtgcgtgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc  
GTGGCGCGAGCTTCTGAAACTTAGGCGGCAGAGCGCGAGCGCGCTGTGGCAGCTGCTGCGCCT  
CTGCTGCGCCTCGGGTGTCTTTTGCGGCGGTGGGTGCGCGCCGGGAGAAGCGTGAGGGGA  
CAGATTTGTGACCGGCGCGGTTTTTGTGAGCTTACTCCGGCCAAAAAAGAACTGCGCCTC  
TGGAGCGGgttagtggtggtggttagtgggttgggacgagcgcgtcttccgcagtcaccagt  
ccagcgtggcgggggagcgccctcagccccgggtcgctgcgcgcgcgcgcgcgcgcgcgcgc  
tctctgccaaacccccaccattgcctgagagaaaggctccttgcgcgaaggcaaattttcgc  
caagcaaattcgagccccgccttccctgggtctccatttccgcctccggcccgccgcct  
ttgggctccgccttcagctcaagacttaacttccctcccagctgtccagatgacgccat  
ctgaaatttcttggaaacacgatcactttaacggaatattgctggtttggggaagtgttt  
tacagctgctgggcaagctgtatttgccctacttaagcccttggaattgctgtattccg  
aagacatgctgatgggaattaccaggcggcgttgggtctctaaactgagccctctgtccca  
actagctacgcgtcactggttagcctga.

## Exon 2

.gacttaactccctcccagctgtcccagatgacgccatctgaaatttcttggaaacacg  
atcactttaacggaatattgctgttttggggaagtgtttacagctgctgggcacgctgt  
atttgccctacttaagccctggtaattgctgtattccgaagacatgctgatgggaatta  
ccaggcgggcgttgggtctctaaactgagccctctgtcccaactagctacgcgtcactggtt

# Fig.7. (2/22) <sup>19/52</sup>

agcgtgattgaaactaaatcgatatgaaaatcctcttctctagtcgcactagccacgtttc  
gagtgccttaatgtggctagtgccaccggtttggacagcacagctgtataaatgttcccatc  
ctcacagtaagctgttaccgttccaggagatgggactgaattagaattcaaacaatttt  
ccagcgttctgagttttacctcagtcacataataaggaatgcatccctgtgttaagtga  
ttttggtcttctgtttttgcagACTTATTTACCAAGCATTGGAGGAATATCGTAGGTAAAA

ATGCCTATTGGATCCAAAGAGAGGCCAACATTTTTTGAAATTTTAAAGACACGCTGCAAC  
M P I G S K E R P T F F E I F K T R C N

20

AAAGCAGgtattgacaaattttatataactttataaattacaccgagaaagtgttttcta  
K A D

23

aaaaatgcttgctaaaaaaccagtcacgtcagtggttgccttagaaccataaaactgttcc  
tatgtgtgtataaatccagttaacaacataatcatcggttgcaggttaaccacatgataa  
atatagaacgtctagtggtataaagaggaaactggcccttgactagcagtaggtaacaat  
tactaacaatcagaagcattaatgttactttatggcagaagtgtccaaactttttggtt  
tcagtactccttatactcttaaaaaatgatctaggacccccggagtgcttttgtttatgta  
gcttaccatattagaaattttaaactaagaatttaaggetgggctgggtggctcacgcct  
gtaatcccagcactttgggaggcccgagggtgggcgatcacttgaggccagaagttag  
accagcctggccaacatggtgaaacc.....

## Exon 3

.....gactacaggcgtgtgccacc  
acgcttgggctaattttttgtgttttagtaaagatgggggttcaacgtgttagcaaggtt  
ggtctcgatctgacctcgtgatctgctcgcctcagcctcccaaagtgttgggattacagg  
cgtgagccccgcacctggccgaattttatcgtggaatgtattcttaatgtgaatagttt  
ttgattccgaacatgaataataagaaaaataaaaaatttaaatgaaaaataaaagctaa  
tatatacagcttttaataatatagttaaatgccatcttgtaacttttgtgaactcttgtt  
acacctttctatagattcgcgaagagaatggattaatgatcttgtttaattaatatgcctt  
aacaaaagtaatccatagtcagatcttaagcatttttttcttatgatctttaactgtt  
ctgggtcacaaaatttgcctgctcactgggttaaaactaagggtgggatttttttttaaatag

ATTTAGGACCAATAAGTCTTAATTGGTTTGAAGAAGCTTTCTTCAGAAGCTCCACCCTATA  
L G P I S L N W F E E L S S E A P P Y N

43

ATTCTGAACCTGCAGAAGAATCTGAACATAAAAACAACAATTACGAACCAAACCTATTTA  
S E P A E E S E H K N N N Y E P N L F K

63

AAACTCCACAAAGGAAACCATCTTATAATCAGCTGGCTTCAACTCCAATAATATTCAAAG  
T P Q R K P S Y N Q L A S T P I I F K E

83

AGCAAGGGCTGACTCTGCCGCTGTACCAATCTCCTGTAAAAGAATTAGATAAATTCAAAT  
Q G L T L P L Y Q S P V K E L D K F K L

103

TAGACTTAGgtaagtaatgcaatatggttagactggggagaactacaaactaggaatttag  
D L G

106

gcaaacctgtgttaaaatcttagctcattcattaattgtgtcatgctgggcaaatcagtc  
tctctggcctcttttctcactcgaaaaatggagacgatgaaaaataatgtctcataggt  
ttggattaaattaaataatgtaggtacttagtaaatgttctctttcatccctcctttgat  
aaatttgccaactgagatttgcgtgaattacgtctttcttatgcaaaaaaacctaggact  
tgttttgatgttaattaaactaaactatatttctgcaagctatcacagaggacagagatt  
attttaccgatatactataagtatcatgatttgggaaggagtttccctggcgtaggtgccg

SUBSTITUTE SHEET (RULE 26)

## Fig.7. (3/22)

20/52

catgttttctaagcaattatgtaataagattatatattcagtcattcaaataattattacc  
tacttgacataagtaataagactttccctt.....

## Exon 4

.....tccttttttagggggaatcagcaaaactgaaaaacctcttcttaca  
actccctatacattctcattcccagtatagaggagactttttgtttttaaacacttccaa  
agaatgcaaatattataatccagagtataacattctcactgaattattgtactgtttcag

GAAGGAATGTTCCCAATAGTAGACATAAAAGTCTTCGCACAGTGAAAACATAAATGGATC  
R N V P N S R H K S L R T V K T K M D Q 126

AAGCAGATGATGTTTCTGTCCACTTCTAAATTCTTGTCTTAGTGAAAGgtatgatgaag  
A D D V S C P L L N S C L S E S 142

ctattatattaaaaatatttaaatgaaacatttttccatcatatatttgttctataaagatg  
aatctgatttttatgctaataattttggcctaagagcctggtagaagatcttacatttttaa  
ataatcttttaggttgagtccttaataagaatagtttttacattagaaacatgtaagttg  
ttgttcttggatgttggaattggctgggttttctgtatattctgtgattttttaagtaaca  
aaaataacagtggtgaaaagcagtaagtcagtccttgaattatcaatttaaaataaattg  
tgtacttttcatctttggagagaatatgatttactttacaaattttttttgttttttt  
tttttttgagatggagtcctgtgcacccaggctgtagtgcagtggtgcgatctcagctca  
ctgcaagctccgcctcccggttcacgccattctcctgcctcagcctcccaagtagctgg  
gactacagg.....

## Exon 5

.....tgcgatctcagctcactgca  
agctccgcctcccggttcacgccattctcctgcctcagcctcccaagtagctgggacta  
caggcgcgccgccaccatgcccggttaattttttgtatttttagtagagacgggtttcac  
tgtgttagctaggatgggtctcgatttctcgacctcgtgatccgcgccctcagcctccca  
gactgctgggattacaggcgtgaaccactgtgcccggcctactttacaaaatttttgagt  
ttaaataacacgggtttccagcagctgaaattttgtgagtacatatgtgttggcattttaaa  
catcacttgatgattatttaatgcttcatgagagatttactttttaaaatgtaataaaa  
atatctaaaagtagtattccaacaatttatatgaatgagaatcttcttttaaaaataaga  
taaactagtttttgccagttttttaaaataacctaagggtatttgctttgttttatttttag

TCCTGTTGTTCTACAATGTACACATGTAACACCACAAAGAGATAAGTCAGgtatgattaa  
P V V L Q C T H V T P Q R D K S V 159

aaacaatgctttttattcttagaataactagaaatgttaataaaaaataaaaacttaacaatt  
ttccctttttttacccccagtggtatgtgggagtttgtttcatacaccaaagtttgtga  
aggtaaatattctacctgggtttatttttargacttagtaattgagaatttgacaatagcg  
ttataacctttgccttgagatttacaaatctgtacctagcattctgcctcatacaggcaat  
tcagtaaacgttaagtgaataaagagtgaaatgaaaaataatattccttaatgatcaggg  
catttctataaaaaataaaactattttcttctccagggctcgtcagacaccacaaacata  
tttctgaaagtctaggagctgaggtggatcctgatatgtcttggtcaagttcttagcta  
caccacccacccttagttctactgtgctcataggtaataatagcaaatgtgtattttacaa  
gaaagagcag.....

## Exon 6

.....gtgttagctaggatgggtctc

Fig.7. (4/22) <sup>21/52</sup>

gatttcctgacctcgtgatccgccgcctcagcctcccagactgctgggattacaggcgt  
gaaccactgtgcccggcctactttacaaaatttttgagtttaaaatacacagggttccagc  
agctgaaatttgtgagtacatatgtgttggcatttttaaacatcacttgatgattatttaa  
tgcttcctgagagatttacttttttaaaatgtaatatataaaatctctaaaagtagtattcca  
acaatttatatgaatgagaatcttcttttaaaaataagataaaactagtttttgccagttt  
tttaaaaataacctaagggatttgccttggttttatttttagtcctgttggcttacaatgtac  
acatgtaacaccacaaagagataagtcagggtatgattaaaaacaatgctttttattctta  
gaatactagaaatgttaataaaaaataaaacttaacaattttccctttttttacccccag

TGGTATGTGGGAGTTTGTTCATACACCAAAGTTTGTGAAGgtaaatattctacctgggt  
V C G S L F H T P K F V K

172

tatttttatgacttagtaattgagaatttgacaatagcgttatacctttgccctgagatt  
tacaaatctgtacctagcattctgcctcatcacaggcaattcagtaaacgttaagtgaat  
aaagagtgaaatgaaaaataaatatccttaatgatcagggcatttctataaaaaataaact  
attttcttttccctccagggctcgtcagacacccaaaacatatcttgaaagtctaggagctg  
agggtggatcctgatatgtcttgggtcaagttcttttagctacaccacccacccttagttcta  
ctgtgctcataggttaataatagcaaatgtgtatttacaagaaagagcagatgaggttgat  
aattgtcatctctaatacttctgtttaaaggaaatatgaaaagaaaatattagataatgt  
ctttgataagtggttagtaactgacaataattttattctattaagtgtagattggaata  
a.....

## Exon 7

.....ccaacaatttatatgaatga  
gaatcttcttttaaaaataagataaaactagtttttgccagtttttttaaaaataacctaagg  
gatttgcttggttttatttttagtcctgttggcttacaatgtacacatgtaacaccacaaa  
gagataagtcagggtatgattaaaaacaatgcctttttattcttagaatactagaaatggtta  
ataaaaaataaaacttaacaattttccctttttttacccccagtggtatgtgggagtttg  
tttcatacaccaaagtttgtgaaggtaaatattctacctgggtttatttttatgacttagt  
aattgagaatttgacaatagcgttatacctttgccctgagatttacaaatctgtacctag  
cattctgcctcatcacaggcaattcagtaaacgttaagtgaataaaagagtgaaatgaaaaa  
ataatatccttaatgatcagggcatttctataaaaaataaactattttcttctcccccag

GGTCGTCAGACACCAAAACATATTTCTGAAAGTCTAGGAGCTGAGGTGGATCCTGATATG  
G R Q T P K H I S E S L G A E V D P D M

192

TCTTGGTCAAGTTCTTTAGCTACACCACCCACCCTTAGTTCTACTGTGCTCATAGgtaat  
S W S S S L A T P P T L S S T V L I V

211

aatagcaaatgtgtattttacaagaaagagcagatgaggttgataattgtcatctctaata  
cttctgtttaaaggaaatatgaaaagaaaatattagataatgtctttgataagtggtgta  
gtaactgacaataattttattctattaagtgtagattggaataaaatacaaatacatttag  
tggtagtcaggtgtgtcaagcattatgttttagtacgatgtgattaacgtagaatagct  
tacaaatatctctttactggcctatataagcgtttaagaggcagttttggtgtgactga  
attctttttacaaatgattgtggttaattggggcattaaagcagc.....

## Exon 8

.....accagtattacattttgttt  
attctagcaaaatagcattctgtttttgattcctcttttagctgggagtaagttaaccctat  
tctgttgcttagatgaaataaatatggataaaatcattttgaaaatatgtatttaatatat  
agtatgcctttaggctgtagtgttgcctaaatgaatgctaaagtctccaagcttttagctt

SUBSTITUTE SHEET (RULE 26)

Fig.7. (5/22) <sup>22/52</sup>

ttaagtcataacctcacagcatcatctgactttccaactcattgtggacagtattaccat  
 aaagtaatgatcaccaagccatatcttaccaccttgtgagtagtactaaggaagtaagta  
 tagtttattcactgtgttgattgacctttctaatctatacttaagtaacttgaatcaat  
 tcattttgtttcaaagtgtgcatgtaatacaatagtagatgtgctttttgatgtctgaca  
 aaaaataagtttttgcattctagtataatatacaatacacataaatttttatcttacag

TCAGAAATGAAGAAGCATCTGAAACTGTATTTCTCATGATACTACTGCTgtaagtaaat  
 R N E E A S E T V F P H D T T A

227

atgacattgatttagactgttgaaattgctaacaattttggaatgccttgttaaattattt  
 atcttacatttttaatttcttaactgttaatttatctaaagccttgagaaagtctctaaa  
 cctggctctatatgtgatttttaacttctgtgaaactctgctgtctctctgttaaagttg  
 catatatacaatatataccgtagtccccatttcatggggatatacattccaatatccccca  
 gtgaatgcttgaaaccttagatagtagcgaaccctatatatatataaaaaatgtgtag  
 tatttatatatatatataacctataatctttttttctataagcacataacctgtgataaag  
 ttaattcataaaattaggcacagtaagagattaacaagaactaataaaaaataggacaa  
 ttataacaaaataaccgtaataaaaagttatgtgaatgttgtctctctgtctcaaaaatatct  
 tattgttctg.....

## Exon 9

.....gcactttatggcttcttttt  
 ggcataatttgaattgccagcatcattatacttgtgctttggggccattgttaagtaaaat  
 aagggtagacttgaacacaagcactgtggtaccacaatagccgatctgataaccaagacaa  
 ctactaagtgactaataggtgggtaccatatacagcctggatacgcctggacaaagggatg  
 attcatgtcccaagtgggatggagcaagatgggtgcaagttttttttctccatttccatt  
 ttcttttcttaagatttccacatcctagtgggtgcaagatttcatcacactactcaggatg  
 acacacaattttaaacttactaattgcttacttctggaattttccattaaaaatttttgg  
 acctaggttgattgcagataactgaaatcaccaaaagtgaaccatggataagggggggac  
 tactactatatgtgcattgagagtttttatactagtatttttaactataatttttgcag

AATGTGAAAAGCTATTTTCCAATCATGATGAAAGTCTGAAGAAAAATGATAGATTTATC  
 N V K S Y F S N H D E S L K K N D R F I

247

GCTTCTGTGACAGACAGTGAAAACACAAATCAAAGAGAAGCTGCAAGTCATGgtaagtc  
 A S V T D S E N T N Q R E A A S H G

265

tctgttttagttgaactacaggtttttttgtgtgtgtgttttgatttttttttttggagg  
 tggagtcctgtctgtcaccogtgatctcagtttaccgcaacctctgcctcccgtgctca  
 agcgatcctgcctcagcttgccaagtagctgagattacaagcatgcaccacatgccc  
 ctattgtatttttagtagagatggcatttcccatgttggccaggctggtctcaaatggt  
 cgtgagccaccatgcccagcctgaactactctttttaattggcaccattgaaggattgct  
 cctcttttcttaagagaaaaatatattacctttcttcttcttgactactgaagtagtattt  
 tatctcaaagrataggagtagaaactaacttggtgtgcctgtgatccagctactcagg  
 aggctgaggtgggaggatcgcttaagcccaggcgggtcaaggttgcagtgagctgtgtgtg  
 tgccactgcact.....

## Exon 10

.....tttttgtctg  
 tgtggatcatgtacgtatgtatatgcataatgtaaaatcagatttacccttgttatagg  
 ccacagaattgatttggacatctgttttgataggtcttagaatatttaattgtatatat  
 agtaagattaggtgagttttaattgtgtagaactgctaagaaaaggttttagggattgt

Fig.7. (6/22)

23/52

tgtatgaataaaaggcttttaggttcattggaatcaggggaatcaggctttactagaagaa  
caggagaaggggtgactgaccgaaaaataaaatgccaagtactcagaataaccctttaaa  
tactgatatgtaatatatttagcacattctacataaaactgtttctatgagaaagggtgtgag  
aataatataaaattatatggcttataaaatattaatgtgcttctgtttttatactttaacag  
  
GATTTGGAAAAACATCAGGGAATTCATTTAAAGTAAATAGCTGCAAAGACCACATTGGAA  
F G K T S G N S F K V N S C K D H I G K 285  
  
AGTCAATGCCAAATGTCCTAGAAGATGAAGTATATGAAACAGTTGTAGATACCTCTGAAG  
S M P N V L E D E V Y E T V V D T S E E 305  
  
AAGATAGTTTTTCATTATGTTTTTCTAAATGTAGAACAAAAAATCTACAAAAAGTAAGAA  
D S F S L C F S K C R T K N L Q K V R T 325  
  
CTAGCAAGACTAGGAAAAAAATTTTCCATGAAGCAAACGCTGATGAATGTGAAAAATCTA  
S K T R K K I F H E A N A D E C E K S K 345  
  
AAAACCAAGTGAAAGAAAAATACTCATTGTATCTGAAGTGGAACCAAATGATACTGATC  
N Q V K E K Y S F V S E V E P N D T D P 365  
  
CATTAGATTCAAATGTAGCACATCAGAAGCCCTTTGAGAGTGGAAGTGACAAAATCTCCA  
L D S N V A H Q K P F E S G S D K I S K 385  
  
AGGAAGTTGTACCGTCTTTGGCCTGTGAATGGTCTCAACTAACCCCTTTCAGGTCTAAATG  
E V V P S L A C E W S Q L T L S G L N G 405  
  
GAGCCCAGATGGAGAAAATACCCCTATTGCATATTTCTTCATGTGACCAAATATTTTCAG  
A Q M E K I P L L H I S S C D Q N I S E 425  
  
AAAAAGACCTATTAGACACAGAGAACAAAAGAAAGAAAGATTTTCTTACTTCAGAGAATT  
K D L L D T E N K R K K D F L T S E N S 445  
  
CTTTGCCACGTATTTCTAGCCTACCAAAATCAGAGAAGCCATTAAATGAGGAAACAGTGG  
L P R I S S L P K S E K P L N E E T V V 465  
  
TAAATAAGAGAGATGAAGAGCAGCATCTTGAATCTCATACTGCACTTCTTGCAGTAA  
N K R D E E Q H L E S H T D C I L A V K 485  
  
AGCAGGCAATATCTGGAACCTTCTCCAGTGGCTTCTTCATTTTCAGGGTATCAAAAAGTCTA  
Q A I S G T S P V A S S F Q G I K K S I 505  
  
TATTCAGAATAAGAGAATCACCTAAAGAGACTTTCAATGCAAGTTTTTTCAGGTCATATGA  
F R I P E S P K E T F N A S F S G H M T 525  
  
CTGATCCAAACTTTAAAAAAGAACTGAAGCCTCTGAAAGTGGACTGGAAATACATACTG  
D P N F K K E T E A S E S G L E I H T V 545  
  
TTTGCTCACAGAAGGAGGACTCCTTATGTCCAAATTTAATTGATAATGGAAGCTGGCCAG  
C S Q K E D S L C P N L I D N G S W P A 565  
  
CCACCACCACACAGAATTCTGTAGCTTTGAAGAATGCAGGTTTAATATCCACTTTGAAAA  
T T T Q N S V A L K N A G L I S T L K K 585  
  
AGAAAACAAATAAGTTTATTTATGCTATACATGATGAAACATCTTATAAAGGAAAAAAA

Fig.7. (7/22)

24/52

K T N K F I Y A I H D E T F Y K G K K I 605  
TACCGAAAAGACCAAAAATCAGAACTAATTAAGTTCAGCCCAGTTTGAAGCAAATGCTT  
P K D Q K S E L I N C S A Q F E A N A F 625  
TTGAAGCACCACTTACATTTGCAAATGCTGATTTCAGgtacctctgtctttttttttttgt  
E A P L T F A N A D S G 637  
aaatagtacatatagttttatagatgacgattccttctgtgtttttttctgtcttttaaa  
atcttcataatcttatatttaattcttaggcattcatctgtatacatgattgttttaggtcttt  
aattaccagtgtttagaatcaggtcactcaaacatggttagataagtttgcatagtttggtg  
tatatccatcactcttgagacagttttattttaagttccgggtacatgtgcaggatgtg  
caggtttgttacataagtaaacgtatgccatgttggtttgctgcacctgtcaaccttca  
cctgagtattaagcccagcatgcattagctatttttccctggtgctctctctccccccaca  
caccccccactcctgacagaccctagtgtgtgtgtgttccctccctgtgtccgtgtgttc  
tcattgttcagctcccacttatgagtgagaacatgt.....

## Exon 11

ctgaaaaatagagcatatttaggattctttctgcttttaaatttgacattcagttattttc  
atgtaatttggtgttttgagcactaccttttaattaattttatttttttttttagag  
actgtctcattctgttacctagtctggagtgcactagtgtgatctcagctcaccgtagcc  
tcacctcctgggctcaagcagtccttgccactcaccctcctgagtaactggcaccacag  
gcatacaccaccacacccagctaattttttttttcatagagtcatggtctcactatgtt  
gcccaggctagtctcgaactcctgggctcaagcagtcctcctgcctcagcctccaaaag  
tgctgagattacaggcatgagccactgtgcccacactacctttttaacttagtgaaaa  
atatttagtgaatgtgattgatggtaactttaattttgtcactttgtgtttttatgtttag  
GTTTATTGCATTCTTCTGTGAAAAGAAGCTGTTTCACAGAATGATTCTGAAGAACCAACTT  
L L H S S V K R S C S Q N D S E E P T L 657  
TGTCCTTAAGTAGCTCTTTTGGGACAATTCTGAGGAAATGTTCTAGAAATGAAACATGTT  
S L T S S F G T I L R K C S R N E T C S 677  
CTAATAATACAGTAATCTCTCAGGATCTTGATTATAAAGAAGCAAATGTAATAAGGAAA  
N N T V I S Q D L D Y K E A K C N K E K 697  
AACTACAGTTATTTATTACCCCGAGAAGCTGATTCTCTGTCATGCCTGCAGGAAGGACAGT  
L Q L F I T P E A D S L S C L Q E G Q C 717  
GTGRAAATGATCCAAAAAGCAAAAAGTTTCAGATATAAAGAAGAGGTCTTGGCTGCAG  
E N D P K S K K V S D I K E E V L A A A 737  
CATGTCACCCAGTACAACATTCAAAAGTGAATACAGTGATACTGACTTTCAATCCCAGA  
C H P V Q H S K V E Y S D T D F Q S Q K 757  
AAAATCTTTTATATGATCATGAAAATGCCAGCACTCTTATTTTAACTCCTACTTCCAAGG  
S L L Y D H E N A S T L I L T P T S K D 777  
ATGTTCTGTCAAACCTAGTCATGATTCTAGAGGCAAAGAATCATACAAAATGTCAGACA  
V L S N L V M I S R G K E S Y K M S D K 797  
AGCTCAAAGGTAACAATTATGAATCTGATGTTGAATTAACCAAAAATATTTCCCATGGAAA  
SUBSTITUTE SHEET (RULE 26)

Fig.7. (8/22)

25/52

L K G N N Y E S D V E L T K N I P M E K 817  
AGAATCAAGATGTATGTGCTTTAAATGAAAATTATAAAAACGTTGAGCTGTTGCCACCTG  
N Q D V C A L N E N Y K N V E L L P P E 837  
AAAAATACATGAGAGTAGCATCACCTTCAAGAAAGGTACAATTCAACCAAAACACAAATC  
K Y M R V A S P S R K V Q F N Q N T N L 857  
TAAGAGTAATCCAAAAAATCAAGAAGAACTACTTCAATTTCAAAAATAACTGTCAATC  
R V I Q K N Q E E T T S I S K I T V N P 877  
CAGACTCTGAAGAACTTTTCTCAGACAATGAGAATAATTTTGTCTTCCAAGTAGCTAATG  
D S E E L F S D N E N N F V F Q V A N E 897  
AAAGGAATAATCTTGCTTTAGGAAATACTAAGGAACTTCATGAAACAGACTTGACTTGTG  
R N N L A L G N T K E L H E T D L T C V 917  
TAAACGAACCCATTTTCAAGAACTCTACCATGGTTTTATATGGAGACACAGGTGATAAAC  
N E P I F K N S T M V L Y G D T G D K Q 937  
AAGCAACCCAAGTGTCATTAATAAAGATTTGGTTTTATGTTCTTGCAGAGGAGAACAAAA  
A T Q V S I K K D L V Y V L A E E N K N 957  
ATAGTGTAAGCAGCATATAAAAAATGACTCTAGGTCAAGATTTAAAAATCGGACATCTCCT  
S V K Q H I K M T L G Q D L K S D I S L 977  
TGAATATAGATAAAATACCAGAAAAAATAATGATTACATGAACAAATGGGCAGGACTCT  
N I D K I P E K N N D Y M N K W A G L L 997  
TAGGTCCAATTTCAATCACAGTTTTGGAGGTAGCTTCAGAACAGCTTCAAATAAGGAAA  
G P I S N H S F G G S F R T A S N K E I 1017  
TCAAGCTCTCTGACATAACATTAAGAAGAGCAAAATGTTCTTCAAAGATATTGAAGAAC  
K L S E H N I K K S K M F F K D I E E Q 1037  
AATATCCTACTAGTTTAGCTTGTTGAAATTGTAAATACCTTGGCATTAGATAATCAAA  
Y P T S L A C V E I V N T L A L D N Q K 1057  
AGAAACTGAGCAAGCCTCAGTCAATTAATACTGTATCTGCACATTTACAGAGTAGTGTAG  
K L S K P Q S I N T V S A H L Q S S V V 1077  
TTGTTTCTGATTGTAAAAATAGTCATATAACCCCTCAGATGTTATTTCCAAGCAGGATT  
V S D C K N S H I T P Q M L F S K Q D F 1097  
TTAATTCAAACCATTAATTTAACACCTAGCCAAAAGGCAGAAATTACAGAACTTTCTACTA  
N S N H N L T P S Q K A E I T E L S T I 1117  
TATTAGAAGAATCAGGAAGTCAGTTTGAATTTACTCAGTTTAGAAAACCAAGCTACATAT  
L E E S G S Q F E F T Q F R K P S Y I L 1137  
TGCAGAAGAGTACATTTGAAGTGCCTGAAAACCAGATGACTATCTTAAAGACCACTTCTG  
Q K S T F E V P E N Q M T I L K T T S E 1157  
AGGAATGCAGAGATGCTGATCTTCATGTCATAATGAATGCCCCATCGATTGGTCAGGTAG



Fig.7. (9/22)

26/52

E C R D A D L H V I M N A P S I G Q V D 1177  
ACAGCAGCAAGCAATTTGAAGGTACAGTTGAAATTAAACGGAAGTTTGCTGGCCTGTTGA  
S S K Q F E G T V E I K R K F A G L L K 1197  
AAAATGACTGTAACAAAAGTGCTTCTGGTTATTTAACAGATGAAAATGAAGTGGGGTTTA  
N D C N K S A S G Y L T D E N E V G F R 1217  
GGGGCTTTTATTCTGCTCATGGCACAAAACCTGAATGTTTCTACTGAAGCTCTGCAAAAAG  
G F Y S A H G T K L N V S T E A L Q K A 1237  
CTGTGAAACTGTTTAGTGATATTGAGAATATTAGTGAGGAACTTCTGCAGAGGTACATC  
V K L F S D I E N I S E E T S A E V H P 1257  
CAATAAGTTTATCTTCAAGTAAATGTCATGATTCTGTGCGTTTCAATGTTTAAGATAGAAA  
I S L S S S K C H D S V V S M F K I E N 1277  
ATCATAATGATAAACTGTAAGTGAAAAAATAATAAATGCCAACTGATATTACAAAATA  
H N D K T V S E K N N K C Q L I L Q N N 1297  
ATATTGAAATGACTACTGGCACTTTTGTGTAAGAAATTACTGAAAATTACAAGAGAAATA  
I E M T T G T F V E E I T E N Y K R N T 1317  
CTGAAAATGAAGATAACAAATATACTGCTGCCAGTAGAAATTCTCATAACTTAGAATTTG  
E N E D N K Y T A A S R N S H N L E F D 1337  
ATGGCAGTGATTCAAGTAAAAATGATACTGTTTGTATTTCATAAAGATGAAACGGACTTGC  
G S D S S K N D T V C I H K D E T D L L 1357  
TATTTACTGATCAGCACAAACATATGTCTTAAATTATCTGGCCAGTTTATGAAGGAGGGAA  
F T D Q H N I C L K L S G Q F M K E G N 1377  
ACACTCAGATTAAAGAAGATTTGTCAGATTTAACTTTTTTGGAAAGTTGCGAAAGCTCAAG  
T Q I K E D L S D L T F L E V A K A Q E 1397  
AAGCATGTCATGGTAATACTTCAAATAAAGAACAGTTAACTGCTACTAAAACGGAGCAAA  
A C H G N T S N K E Q L T A T K T E Q N 1417  
ATATAAAAGATTTTGAGACTTCTGATACATTTTTTTCAGACTGCAAGTGGGAAAAATATTA  
I K D F E T S D T F F Q T A S G K N I S 1437  
GTGTCGCCAAAGAGTCATTTAATAAAATTGTAAATTTCTTTGATCAGAAACCAGAAGAAT  
V A K E L F N K I V N F F D Q K P E E L 1457  
TGCATAACTTTTCCTTAAATTCTGAATTACATTCTGACATAAGAAAGAACAAAATGGACA  
H N F S L N S E L H S D I R K N K M D I 1477  
TTCTAAGTTATGAGGAAACAGACATAGTTAAACACAAAATACTGAAAGAAAGTGTCCCAG  
L S Y E E T D I V K H K I L K E S V P V 1497  
TTGGTACTGGAATCAACTAGTGACCTTCCAGGGACAACCCGAACGTGATGAAAAGATCA  
G T G N Q L V T F Q G Q P E R D E K I K 1517  
AAGAACCTACTCTGTTGGGTTTTTCATACAGCTAGCGGGAAAAAAGTTAAAAATTGCAAAGG

Fig.7. (10/22)

27/52

E P T L L G F H T A S G K K V K I A K E 1537  
AATCTTTGGACAAAGTGAAAAACCTTTTTGATGAAAAAGAGCAAGGTACTAGTGAAATCA  
S L D K V K N L F D E K E Q G T S E I T 1557  
CCAGTTTTAGCCATCAATGGGCAAAGACCCTAAAGTACAGAGAGGCCTGTAAAGACCTTG  
S F S H Q W A K T L K Y R E A C K D L E 1577  
AATTAGCATGTGAGACCATTGAGATCACAGCTGCCCCAAAGTGTAAGAAATGCAGAATT  
L A C E T I E I T A A P K C K E M Q N S 1597  
CTCTCAATAATGATAAAAAACCTTGTTTCTATTGAGACTGTGGTGCCACCTAAGCTCTTAA  
L N N D K N L V S I E T V V P P K L L S 1617  
GTGATAATTTATGTAGACAAACTGAAAATCTCAAAACATCAAAAAGTATCTTTTTGAAAG  
D N L C R Q T E N L K T S K S I F L K V 1637  
TTAAAGTACATGAAATGTAGAAAAAGAAACAGCAAAAAGTCCTGCAACTTGTTACACAA  
K V H E N V E K E T A K S P A T C Y T N 1657  
ATCAGTCCCCTTATTCAGTCATTGAAAATTCAGCCTTAGCTTTTTACACAAGTTGTAGTA  
Q S P Y S V I E N S A L A F Y T S C S R 1677  
GAAAAACTTCTGTGAGTCAGACTTCATTACTTGAAGCAAAAAAATGGCTTAGAGAAGGAA  
K T S V S Q T S L L E A K K W L R E G I 1697  
TATTTGATGGTCAACCAGAAAGAATAAATACTGCAGATTATGTAGGAAATTATTTGTATG  
F D G Q P E R I N T A D Y V G N Y L Y E 1717  
AAAATAATTCAAACAGTACTATAGCTGAAAATGACAAAAATCATCTCTCCGAAAAACAAG  
N N S N S T I A E N D K N H L S E K Q D 1737  
ATACTTATTTAAGTAACAGTAGCATGTCTAACAGCTATTCCTACCATTCTGATGAGGTAT  
T Y L S N S S M S N S Y S Y H S D E V Y 1757  
ATAATGATTCCAGGATACTCTCAAAAAATAAATTGGATTCTGGTATTGAGCCAGTATTGA  
N D S R I L S K N K L D S G I E P V L K 1777  
AGAATGTTGAAGATCAAAAAACACTAGTTTTTCCAAAGTAATATCCAATGTAAAAGATG  
N V E D Q K N T S F S K V I S N V K D A 1797  
CAAATGCATACCCACAAACTGTAAATGAAGATATTTGCGTTGAGGAACTTGTGACTAGCT  
N A Y P Q T V N E D I C V E E L V T S S 1817  
CTTCAACCCTGCAAAAAATAAAATGCAGCCATTAAATTGTCCATATCTAATAGTAATAATT  
S P C K N K N A A I K L S I S N S N N F 1837  
TTGAGGTAGGGCCACCTGCATTTAGGATAGCCAGTGGTAAAATCGTTTGTGTTTCACATG  
E V G P P A F R I A S G K I R L C S H E 1857  
AAACAATTA AAAAAGTGAAAGACATATTTACAGACAGTTTCAGTAAAGTAATTAAGGAAA  
T I K K V K D I D S F S K V I F T K E N 1877  
ACAACGAGAATAAATCAAAAATTTGCCAAACGAAAATTATGGCAGGTTGTTACGAGGCAT

Fig.7. (11/22)

28/52

N E N K S K I C Q T K I M A G C Y E A L 1897  
TGGATGATTTCAGAGGATATTCTTCATAACTCTCTAGATAATGATGAATGTAGCACGCATT  
D D S E D I L H N S L D N D E C S M H S 1917  
CACATAAGGTTTTTCTGACATTTCAGAGTGAAGAAATTTTACAACATAACCAAAATATGT  
H K V F A D I Q S E E I L Q H N Q N M S 1937  
CTGGATTGGAGAAAGTTTCTAAAATATCACCTTGTGATGTTAGTTTGGAAACTTCAGATA  
G L E K V S K I S P C D V S L E T S D I 1957  
TATGTAAATGTAGTATAGGGAAGCTTCATAAGTCAGTCTCATCTGCAAATACTTGTGGGA  
C K C S I G K L H K S V S S A N T C G I 1977  
TTTTTAGCACAGCAAGTGGAAAATCTGTCCAGGTATCAGATGCTTCATTACAAAACGCAA  
F S T A S G K S V Q V S D A S L Q N A R 1997  
GACAAAGTGTCTGAAATAGAAGATAGTACCAAGCAAGTCTTTTCCAAAGTATTGTTTA  
Q V F S E I E D S T K Q V F S K V L F K 2017  
AAAGTAACGAACATTTCAGACCAGCTCACAAGAGAAGAAAATACTGCTATACGTACTCCAG  
S N E H S D Q L T R E E N T A I R T P E 2037  
AACATTTAATATCCCAAAAAGGCTTTTCATATAATGTGGTAAATTCATCTGCTTTCTCTG  
H L I S Q K G F S Y N V V N S S A F S G 2057  
GATTTAGTACAGCAAGTGGAAAGCAAGTTTCCATTTTAGAAAGTTCCTTACACAAAGTTA  
F S T A S G K Q V S I L E S S L H K V K 2077  
AGGGAGTGTAGAGGAATTTGATTTAATCAGAACTGAGCATAGTCTTCACTATTACCTA  
G V L E E F D L I R T E H S L H Y S P T 2097  
CGTCTAGACAAATGTATCAAAAATACTTCCTCGTGTGATAAGAGAAACCCAGAGCACT  
S R Q N V S K I L P R V D K R N P E H C 2117  
GTGTAAACTCAGAAATGGAAAAACCTGCAGTAAAGAATTTAAATTATCAAATAACTTAA  
V N S E M E K T C S K E F K L S N N L N 2137  
ATGTTGAAGGTGGTTCTTCAGAAAATAATCACTCTATTAAAGTTTCTCCATATCTCTCTC  
V E G G S S E N N H S I K V S P Y L S Q 2157  
AATTTCAACAAGACAAACAACAGTTGGTATTAGGAACCAAAGTCTCACTTGTGAGAACA  
F Q Q D K Q Q L V L G T K V S L V E N I 2177  
TTCATGTTTTGGGAAAAGAACAGGCTTCACCTAAAAACGTAAAAATGGAAATTGGTAAAA  
H V L G K E Q A S P K N V K M E I G K T 2197  
CTGAAACTTTTTCTGATGTTCTGTGAAAACAAATATAGAAGTTTGTCTACTTACTCCA  
E T F S D V P V K T N I E V C S T Y S K 2217  
AAGATTTCAGAAACTACTTTGAAACAGAAGCAGTAGAAATTGCTAAAGCTTTTATGGAAG  
D S E N Y F E T E A V E I A K A F M E D 2237  
ATGATGAACTGACAGATTCTAAACTGCCAAGTCATGCCACACATTCTCTTTTACATGTC

SUBSTITUTE SHEET (RULE 26)

Fig.7. (12/22) <sup>29/52</sup>

D E L T D S K L P S H A T H S L F T C P 2257  
CCGAAATGAGGNAATGGTNTTGTCAAATTCAGAATTGGAAAAAGAAGAGGAGAGCCCC  
E N E E M V L S N S R I G K R R G E P L 2277  
TTATCTTAGTGGGtaagtggttcatttttacctttcgtgttgccaatcactatttttaag  
I L V G 2281

tgtttattcagtagacttggtatgctaacaattaagagtggtataaaactatgtcttttca  
gccatttttgtgtagtcagtttgggggagtatggtttgatatacagatacacagattcag  
tattcgtatacagatttgatatcttgggtatacagattcgatatctctgaatctgtatacc  
aaqaaatcatgttttaagggtctcaatatattttcaaaaagattattagtataaataattq  
agaaattactgttaaaaagttttgagtttctctagaaaatttgaaactcttaacaaaacc  
tgcataataacttaactgttttcatatacatagcaagttcagactctgacttatatg  
aactttaaaagttggtttccggggaggccgaggcgggaggatcacgaggtcaggagatcga  
gaccatcccggc.....

## Exon 12

gtaatttgataattgtaaaaagcctcttaactctaattcaaggacctacataataaatta  
ctccttcagttaatggctgcccccgctgctgaaaaaagagagaaaaagttt  
atttgaagaaattttgttaggccttattgccagtaaacctagagttatatattagtgtcag  
tttttcaaaaagtagcttatctgtggtatctggttagcatctgtttatcctatttaggatt  
tatcctgttttagacctgttaaatagtggtgttttaagtggtcaaaacagaaacaaaat  
gtaattgacattgaagactgactttactctttcaaacattaggtcactatttgttgtaag  
tatttttgtttaacatttaagaggtcaatacttttagctttaaaaaaatggtctatagact  
tttgagaaataaaactgatatttttgccttaaaaacatatatgaaatatttcttttttag

GAGAACCCTCAATCAAAAAGAACTTATTAATGAATTTGACAGGATAATAGAAAATCAAG  
E P S I K R N L L N E F D R I I E N Q E 2301

AAAAATCCTTAAGGCTTCAAAAAGCACTCCAGATGgttaaaattagctttntattnta  
K S L K A S K S T P D G 2313

tctgtttctccctctataggttatgggtatataatattctgacctcaggtggatccacctg  
cctctccaaagtgtggtggnattacagacatgagccactgtgcctaatacaggacctcttt  
atactcttaaaaattactgaggacctaaagnggcatttgtttatgtgggantatatctat  
tgatatttaccatattnganatgtaaattgattaatggttaaaanttagtaantattatgc  
gttggtcattgggaggatattgagttcactgagttatgcggatcttccgaaagttgaacag  
ttctattatgccagtaattaaaacaatccacctttccattggatgccattaccggatc  
cagaaaagnttaaagtagtagaaagctgtcaagcttacagagcccagatacaagcttccc  
caaaaattctgattttcatctaaaagcttgaatttt.....

## Exon 13

gagatcgctcctcattcttttttgtggttacatagtagttgatcatctggctgtgtcagtg  
tttcttagttttatttaaccaattttccaactagtggtacttattgaagatttaattaggttc  
cagttacatactgaaaatgaacaatatctaaagcttagcttttaaaccttcataagacta  
aattttaaatttgggtatttgcatacagaatttagctaacacctttgagttatgatggttaa  
catcaactgactaaatttatgctgattttctgttggtatgcttgtaactgtgagttatttggg  
gcatagtcattatcaatttgtgaatcaatttattttcatagtttaacatttattgagcatc  
cgttacattcactgaaaattgtaaagcctataattgtctcaaattttttgtgtatttaca  
gtaacatggatattctcttagattttaactaatatgtaataataaaataattgttttcttag

SUBSTITUTE SHEET (RULE 26)

30/52

TACCCTTTCGgtaagacatgtttaaatttttctaaattctaatacagtatgagaaaagtc  
P E R 2336

Exon 14

CACAACTAAGGAACGTCAAGAGATACAGAATCCAAATTTTACCGCACCTGGTCAAGAATT  
T T K E R Q E I Q N P N F T A P G Q E F 2356

TCTGTCTAAATCTCATTTGTATGAACATCTGACTTTGGAAAAATCTTCAAGCAATTTAGC  
L S K S H L Y E H L T L E K S S S N L A 2376

AGTTTCAGGACATCCATTTTATCAAGTTTCTGCTACAAGAAATGAAAAAATGAGACACTT  
V S G H P F Y O V S A T R N E K M R H L 2396

GATTACTACAGGCAGACCAACCAAAGTCTTTGTTCCACCTTTTAAAACTAAATCACATTT  
I T T G R P T K V F V P P F K T K S H F 2416

TCACAGAGTTGAACAGTGTGTTAGGAATATTAAC TTGGAGGAAAAACAGACAAAAGCAAAA  
H R V E Q C V R N I N L E E N R Q K Q N 2436

CATTGATGGACATGGCTCTGATGATAGTAAAAATAAGATTAATGACAATGAGATTCATCA  
I D G H G S D D S K N K I N D N E I H Q 2456

GTTTAACAAAAACAACCTCCAATCAAGCAGCAGCTGTAACCTTTACAAAGTGTGAAGAAGA  
 F N K N N S N O A A A V T F T K C E E E 2476

ACCTTTAGgtatcgatatgacaatttgtgtgatgaatttttgcctttcagttagatatattc  
P L D 2479

cggtgttaaataatgtcctgatgggttttccccctttgggtgggtggaattttaaagccctt  
tttaatgttttagattttctaaatccaaagattaggtttaaattattctaatgtttcttt  
caaagataactcttgtggacttggttaaaaaaaattagacacacaatct.....

Fig.7. (14/22)

3/52

## Exon15

.....aaccactgtgcctgg  
ccaggggttgctgttttaaatccaattttatttttgctaagtatttattccttgatag  
ATTTAATTACAAGTCTTCAGAATGCCAGAGATATACAGGATATGCGAATTAAGAAGAAAC  
L I T S L Q N A R D I Q D M R I K K K Q 2499  
AAAGGCAACGCGTCTTTCCACAGCCAGGCAGTCTGTATCTTGCAAAAACATCCACTCTGC  
R Q R V F P Q P G S L Y L A K T S T L P 2519  
CTCGAATCTCTCTGAAAGCAGCAGTAGGAGGCCAAGTTCCCTCTGCgtgtccccataaac  
R I S L K A A V G G Q V P S A 2534  
aggatgtgttttgnctacaatactgatggcttttatgacagagtgttaattttatttcatt  
aactagtatcctacaaatggctttgttttaagaatgaacacattagtgcaggaatggatg  
aatgaaatcatccatattttcctaattaagccctgcagtgccagcctctggcccttgct  
aggcctgccctcaccctactaaagtgatctgtgccttcccaaattactacttctttcc  
ccccttcaaacttttctattttgtcattgtaaatgctctcagctagggtgttaaagtagt  
cttactgatattcaaagtgaataactgatagccctgaaccttctatgagctatttatat  
ttccaaagaggattctccttaagccaatattatctaggtagaatttttaggcaatggaga  
ggtgaaaataatattgatgacattaatagctaacttttgagcatttt.....

## Exon 16

gtctgactccagagtcaaactctgaacaaacaaaaagacactttgggttagatatcctgg  
ggtgaaagcaagcactttgaaagtaagccaagcctgtgtacagatctgaccacctgaggt  
cacattccctaaaaataactttaaacttctccctttgtttcccatctaagtttttgaactta  
agagattttgtaaaacatcacatttttttatectcacagtaccttccatggcagattta  
gcaggaggcgtataaacggggtggaaaagggtacagcagactgtggaatgtatggatcatt  
tatattacattaaaatttttagtttctagtaaataacttaaagtgtttttagtgaagat  
tctagtagttaatgaaaatttttggtaaattcagttttgggtttgttataattgtttttat  
tgtgtgatacatgtttactttaaattgtttttctttttgtgtgtgtttattttgtgtag  
GTGTTCTCATAAACAGCTGTATACGTATGGCGTTTCTAAACATTGCATAAAAATTAACAG  
C S H K Q L Y T Y G V S K H C I K I N S 2554  
CAAAAATGCAGAGTCTTTTCAGTTTCACACTGAAGATTATTTTGGTAAGGAAAGTTTATG  
K N A E S F Q F H T E D Y F G K E S L W 2574  
GACTGGAAAAGGAATACAGTTGGCTGATGGTGGATGGCTCATACCCCTCCAATGATGGAAA  
T G K G I Q L A D G G W L I P S N D G K 2594  
GGCTGGAAAAGAAGAATTTTATAGgtactctatgcaaaaagattgtgtgttaacttttat  
A G K E E F Y R 2602  
gtattccctcaccctcttttcttctttaaactgtctctcgaactaaaaagttggctagaa  
atcaaatttttatgcatttaattgttttaagtgcattatgggttaagcattctgtagaagt  
cttttgaaaagtgtgtttgtccctggggtttaatgaactggattttcttgatttgggaca  
tttttcttaggcattttataaatatagcccaattttataaagttaaatttggccgggtacag  
tggctcatgcctgtaatcccagcactttgggaggccgagggcgggttagatcacctgaggtc  
aggagttcagagaccagcctggccaacgtggcgaaaccccatctctactaaaagttacaaga

SUBSTITUTE SHEET (RULE 26)

Fig.7. (15/22) <sup>32/52</sup>

actatctgggctggtgggcaggcacctgtaatccccggctactctggaggctgaggcagga  
gaatcgcttgaacctgggaggcag.....

## Exon 17

.....ttgttttgtttttatatattt  
gagatagggtctcactcttgtccaggctggagtgagtgaggcactatcatggctcactgca  
gcctcaacctcctgggctcaagcaatcctcccaccacagcctcctaagtagctgggacca  
cagatgtgagctaccactcttggctgattttttttattattttttgtagagatgtggggg  
tctcactatgttgccctaggctgggtctcaaacttctggcctcaagcaatcctcctgcctca  
gcttcccaaaatgctgggagtataggcatgagccaccatgctcagcaatgaagtttttat  
cagtatgatactttgatacatgtcaaataattttctgaaattatattgtagatcatatga  
actcataaaaacttaatgatcttgaacaatgtagttttttgracagagaatagttgtagtt  
gttgaattcagtatcatcctatgtgggtttttatgataaatattctactttttattgttcag

GGCTCTGTGTGACACTCCAGGTGTGGATCCAAAGCTTATTTCTAGAATTTGGGTTTATAA  
A L C D T P G V D P K L I S R I W V Y N 2622

TCACTATAGATGGATCATATGGAACTGGCAGCTATGGAATGTGCCTTTTCTAAGGAATT  
H Y R W I I W K L A A M E C A F P K E F 2642

TGCTAATAGATGCCTAAGCCCAGAAAGGGTGCTTCTTCAACTAAAATACAGgcaagttaa  
A N R C L S P E R V L L Q L K Y R 2659

aagcattacattacgtaatcatatacggcagtgatgggttaagggtttctgtgtagtctgtg  
acttccatgtcaaaaatgttgacacagccagttgtcagtgacagttgccatcccacactgc  
tgttctcctgtcatccctagccccattttaagagagatcacacattcatgcattgctgct  
ttccctctttccccacccccctccttaacctcttgatgtatgagaagaatatgagttacta  
atttgatccactatttgggggattgctaataaagcattttttgcattttattttttgctttt  
taaaaaataattgatattttaacaatatgaacaatatattcctagctacaaaatttttaa  
ttctcagtatctcttagataaattcagtttttattctcagttattcagtgacttggttaa  
acagtggaattcttagagtcacacttccctaaaatatgcattttttgttttcacttttagata  
tgatacgga.....

## Exon 18

.....caactaaaatacagggaagt  
ttaaagcattacattacgtaatcatatacggcagtgatgggttaagggtttctgtgtagtct  
gtgacttccatgtcaaaaatgttgacacagccagttgtcagtgacagttgccatcccacac  
tgctgttctcctgtcatccctagccccattttaagagagatcacacattcatgcattgct  
tgcttccctctttccccacccccctccttaacctcttgatgtatgagaagaatatgagtta  
ctaatttgatccactatttgggggattgctaataaagcattttttgcattttattttttgct  
ttttaaaaaataattgatattttaacaatatgaacaatatattcctagctacaaaatttt  
taattctcagtatctcttagataaattcagtttttattctcagttattcagtgacttggt  
taaacagtggaattcttagagtcacacttccctaaaatatgcattttttgttttcacttttag

ATATGATACGGAAATTGATAGAAGCAGAAGATCGGCTATAAAAAAGATAATGGAAAGGGA  
Y D T E I D R S R R S A I K K I M E R D 2679

TGACACAGCTGCAAAAACACTTGTCTCTGTGTTTCTGACATAATTTTCATTGAGCGCAAA  
D T A A K T L V L C V S D I I S L S A N 2699

TATATCTGAAACTTCTAGCAATAAACTAGTAGTGCAGATACCCAAAAAGTGCCATTAT  
I S E T S S N K T S S A D T Q K V A I I 2719

Fig.7. (16/22)

33/52

TGAACTTACAGATGGGTGGTATGCTGTTAAGGCCAGTTAGATCCTCCCCTCTTAGCTGT  
E L T D G W Y A V K A Q L D P P L L A V 2739  
CTTAAAGAATGGCAGACTGACAGTTGGTCAGAAGATTATTCTTCATGGAGCAGAAGTGGT  
L K N G R L T V G Q K I I L H G A E L V 2759  
GGGCTCTCCTGATGCCTGTACACCTCTTGAAGCCCCAGAATCTCTTATGTTAAAGGttaa  
G S P D A C T P L E A P E S L M L K 2777

ttaatttgcactcttggtaaaaatcagtcattgattcagttaaattctagaagttttaca  
tttaaatttttaaagcttactaaggatgctcaattttcttagatgtactgataatttttagt  
ataaaaaacatattcttcagacaqttaaagtttttgtgcagtttttgggaggtccagaga  
tctttcttgagcttaaataatgcatttccaattaaaaagcaaaataaatttgcaccattt  
gattttgggtatctgtagcttgcctccttctgttctcatagctttgctttgatcagatcc  
ctattccactctggattagagaattacatttttagtacttttcaaataatgtaatagataca  
ctttttatctctatgtagatttttaaactacataacaggactctttgtcatattgaatggg  
ctgcagttattgctatctgaaattaccgataatattgtacattcagattcacttaagaggt  
aaccttgcagagaat.....

## Exon 19

.....cttaacaccatgccaggta  
ccttagtaagtgttcgatgaataatttgcctttttgtattagccataatcattctcaggctg  
ctttgtcattttacttgttccacaaattcttagcttccaaaattttgggtgatacctcattt  
cctattctctctagttgcctttgtccatgtagattttttgaggaagcttgggtaaaataag  
tgtatttttaaactattatgttttaaactgaagttccttttatctgttttctaatagaaca  
tttaaataagcatttaagaactttagtagcagtataaacaatatgtttgagaagtactatattg  
tgaaaatattttcactttttatacagtttttttacttatttactgtcttactaatcttccta  
agacttttttaagtgaatatttttaaggcagttctagaagaatgaaaactcttatgatat  
ctgtaatagaattgaatacatattttaactactaaatcaatatattttattaatttgtccag

ATTTCTGCTAACAGTACTCGGCCTGCTCGCTGGTATACCAAAGTGGATTCTTTCCTGAC  
I S A N S T R P A R W Y T K L G F F P D 2797

CCTAGACCTTTTCCTCTGCCCTTATCATCGCTTTTCAGTGATGGAGGAAATGTTGGTTGT  
P R P F P L P L S S L F S D G G N V G C 2817

GTTGATGTAATTATTCAAAGAGCATACCTATACAGGtatgatgtattcttgaaacttac  
V D V I I Q R A Y P I Q 2829

catatattttctttcttttgatacaattaatttgtttgtttgtttgagatggagtttcggg  
ctcttggccaggctggagtgcaatggcgatcttggttcactgcagcctcccacctccc  
gggttcaagtgaattctctgcctcagcctctcaagtagctgagccaccacacctgggctaa  
ttttgtatttttggtagagaaggggtttcatcatgttgggtcaggctgatctcgaactcct  
gacctcaggtgatccactaatctcagcctcccaaagttctgggattacagatgtgagcca  
ctgtgcctggcctgatacaattaacttgaatgttatatatgtgacttttttgggtgtgtgt  
aacacattattacagtggtgagagaagacatcatctggattatacatatttcgcaatgaa  
agagaggaagaaaaggaagcagcaaaatatgtggaggcccaacaaaagagactaga....

## Exon 20

.....ctgaccctagaccttttccct  
ctgcccttatcatcgctttttcagtgatggaggaaatgttgggtgtgttgatgtaattatt

SUBSTITUTE SHEET (RULE 26)



Fig.7. (17/22)

34/52

caaagagcataccctatacaggtatgatgtattcttgaaacttaccatatatttctttct  
tttgatacaattaatttgtttgtttgttttgagatggagtttcggtctcttgcccaggctg  
gagtgcaatggcgtgatcttgggttactgcagcctccacctcccggttcaagtgatcc  
tcctgcctcagcctctcaagtagctgagccaccacacctggctaattttgtatttttggg  
agagaaggggtttcatcatgttgggtcaggtgatctcgaactcctgacctcaggtgatcc  
actaatctcagcctcccaaagttctgggattacagatgtgagccactgtgcctggcctga  
tacaattaacttgaatgttatatatgtgacttttttgggtgtgtgtaacacattattacag

TGGATGGAGAAGACATCATCTGGATTATACATATTTTCGCAATGAAAGAGAGGAAGAAAAG  
R M E K T S S G L Y I F R N E R E E E K 2849

GAAGCAGCAAAATATGTGGAGGCCCAACAAAAGAGACTAGAAGCCTTATTCATAAAATT  
E A A K Y V E A Q Q K R L E A L F T K I 2869

CAGGAGGAATTTGAAGAACATGAAGGtaaaattagttatatgtgtacacattgttatttct  
Q E E F E E H E E 2878

aatatgagaacaaagtcttagagactttgaatttaacatttttaagttagtaaaattgtttt  
tattttgagtagtaaaattgactttatttttagtatctagggatttcttttttgggtgta  
gacaaagaatagcaacaaggacagaaatatcaggtctaaagccatttgaatatattttcc  
tgaattcttacctatatgatgtggcttttgcatttttgtcatggtagttattagctttca  
tgtgttattatgcctggaactaggacctattgtgggtgtcaattttaataaaaaaatcat  
ggtgttttgatgtttatatgacataaaattttatttttctgtatctcccttttgtgtgtgc  
tgaagattttatgtttttctgcatttctcatgatttatatagatgtaacatgttctata  
ggacatgtaattttacatgtcctatagaactataagttacatgtcc.....

## Exon 21

.....atncattatacgaagttatggatcacaccactgcactc  
ccagattgggtgacagagtgcagacctgtctcaaaaaaaaaaaaaagaaaaacttttag  
cagttatatagtttcttatctttaaatctcccttctttgggtgttttatgcttggttctt  
tagtttttagttgcttttgaatttacagtttagtgaattaataatccttttgttttcttag

AAAACACAACAAAACCATATTTACCATCACGTGCACTAACAAGACAGCAAGTTCGTGCTT  
N T T K P Y L P S R A L T R Q Q V R A L 2898

TGCAAGATGGTGCAGAGCTTTATGAAGCAGTGAAGAATGCAGCAGACCCAGCTTACCTTG  
Q D G A E L Y E A V K N A A D P A Y L E 2918

AGgtgagagagtaagaggacatataatgaggcttgatgattattcaaggtgagaagctgt  
tttanactctctggccatcacaggaaggaatatgttgaaatgctgcatttctccaaaagg  
gatntgttcatttctgggattttccagtgatgttgccagac.....

## Exon 22

.....gtatctactatttcaaagtt  
aatggaattatactcctggggctaagaatgagggttctagggccaacctctactacctat  
gtggcttgtgcaaattagttgtccctttgtgcctcagttttacctacaacacagaaaca  
atgatattacctaccccatggactgttgtgaagattaaatgaattagtagacatttactaca  
catagatctatttctcaaaataatgagcattcagatattagccatctgtaattgtagtgg  
tgatgattatgattattagagtacatttataattggaggatcatttttgcctgtagggaaa  
tagaattattaatagtttggagcacctgagaatattatgtgagaaactgattacattaac  
cacaccttaagatgagctcctaattttgttggattttgtcctgttttaaagccatctagtta  
caatagatggaacttttttgttctgattgtctttttattccaatatcttaaatgggtcacag

35/52

Fig.7. (18/22)

GGTTATTTTCAGTGAAGAGCAGTTAAGAGCCTTGAATAATCACAGGCAAATGTTGAATGAT  
G Y F S E E Q L R A L N N H R Q M L N D 2938

AAGAAACAAGCTCAGATCCAGTTGGAAATTAGGAAGGCCATGGAATCTGCTGAACAAAAG  
K K Q A Q I Q L E I R K A M E S A E Q K 2958

GAACAAGGTTTATCAAGGGATGTCACAACCGTGTGGAAGTTGCGTATTGTAAGCTATTCA  
E Q G L S R D V T T V W K L R I V S Y S 2978

AAAAAAGAAAAAGATTTCAGGtaagtatgtaaagctttgtttttatcagttttattaact  
K K E K D S V 2985

taaaaaatgaccttactaacaacaaatgattataaatccagataaagtataaagttagttta  
tatcagagaagcaaaatccactactaatgcccacaaagagataatataaaagaggatctg  
tatttttttgaacaaacattttaaagataatcacttcttccattgcatctttctcatc  
tttctccaaacagttatactgagtatttggcgtccatcatcagatttatattctctgtta  
acagaaggaaagagatacagaatttatcatcttgcaacttcaaaatctaaaagtaaatct  
gaaagagctaacatacagtttagcagcgacaaaaaaactcagtatcaacaactaccggta  
caaacctttcattgttaatttttgcagttttgataagtgcttgtagtttatggaatctcca  
tatgttgaatttttgttttgttttctgtaggtttcagat.....

## Exon 23

.....ctagttacaatagatggaac  
ttttttgttctgattgctttttattccaatatcttaaaggtcacagggttatttccagt  
aagagcagtttaagagccttgaataatcacaggcaaatgttgaatgataagaaacaagctc  
agatccagttggaaatttaggaaggccatggaatctgctgaacaaaaggaacaaggttat  
caagggatgtcacaaccgtgtggaagttgcgtattgtaagctattcaaaaaaagaaaaag  
attcaggtaagtatgtaaagctttgtttttatcagttttattaacttaaaaaatgacct  
tactaacaacaaatgattataaatccagataaagtataaagttagtttatatcagagaagca  
aatccactactaatgcccacaaagagataatataaaagaggatctgtattttttgaa  
acaaacattttaaagataatcacttcttccattgcatctttctcatctttctccaaacag

TTATACTGAGTATTTGGCGTCCATCATCAGATTTATATTCTCTGTTAACAGAAGGAAAGA  
I L S I W R P S S D L Y S L L T E G K R 3005

GATACAGAATTTATCATCTTGCAACTTCAAAATCTAAAAGTAAATCTGAAAGAGCTAACA  
Y R I Y H L A T S K S K S K S E R A N I 3025

TACAGTTAGCAGCGACAAAAAACTCAGTATCAACAACCTACCGgtacaaacctttcatt  
Q L A A T K K T Q Y Q Q L P 3039

gtaatttttgcagttttgataagtgcttgtagtttatggaatctccatagttgaaat  
tggtttgttttctgtaggtttcagatgaaatttatttccagatttaccagccacgggagc  
cccttcacttcagcaaattttagatccagactttcagccatcttggttctgaggtggacc  
taataggatttgcgtttctggttgtaaaaaaacaggtaatgcacaatatagtttaatttt  
ttttattgattcttttaaaaaacattgtcttttaaaatctcttatgattagttggagcta  
ccagttggcaaatttgctagctaaactagtgatctgaaagtaagcctctttgaacctctga  
tttttcatgaaaagcaattctctcaattctatatttttcaagggttaacaagttacatcc  
tagtctgtgtacttaattttatagaaattgtccttaattttattttctgcaatttatgtt  
ttct.....

Exon 24

Fig.7. (19/22)

36/52

.....aagattcaggtaagtatgta  
aatgctttgtttttatcagttttattaacttaaaaaatgaccttactaacaaaatgatta  
taaataccagataaagtataaaagtttagttttatcagagaagcaaaaatccactactaatgc  
ccacaaagagataataataaaagaggatctgtattttttgaaacaaacatttaaagat  
aatcacttcttccattgcatctttctcatctttctccaaacagttatactgagtatttgg  
cgtccatcatcagatttatattctctgtttaacagaaggaaagagatacagaatttatcat  
cttgcaacttcaaaaatctaaaagtaaatctgaaagagctaacatacagtttagcagcgaca  
aaaaaaactcagtatcaacaactaccggtacaaacctttcattgtaatttttcagtttgg  
ataagtgcttgtagtttatggaatctccatagtgtgaatttttgttttgttttctgtag

GTTTCAGATGAAATTTTATTTTCAGATTTACCAGCCACGGGAGCCCCCTTCACTTCAGCAAA  
V S D E I L F Q I Y Q P R E P L H F S K 3059

TTTTTAGATCCAGACTTTCAGCCATCTTGTCTGAGGTGGACCTAATAGGATTTGTCGTT  
F L D P D F Q P S C S E V D L I G F V V 3079

TCTGTTGTGAAAAAACAGgtaatgcacaatatagttaattttttttattgattctttta  
S V V K K T G 3086

aaaaacattgtcttttaaaatctcttatgattagttggagctaccagttggcaaatttgc  
tagctaactagtgtatctgaaagtaagcctctttgaacctctgatttttcatgaaaagcaa  
ttctctcaattctatatattttcaagggtacaagttacatcctagtctgtgtacttaat  
tttatagaaattgtccttaattttattttctgcaatttatgttttcttactatttctggt  
gtatgtgtttatcccattgtgatgttatattgggtgtcctcaatttatttcccttagccata  
cactctacttttcatgtacagggctattttattatctcagagtcgaagctttttttttt  
tttttttccccgagatggagctctcactctgttggccaggctggagtcagtgggcnaat  
ntcagccccacngcaagtctctgctcccagggttnanacca.....

Exon 25

.....tgtatatgagctaataaaag  
ttgcttccaataattttataattataatgttttcagtgagtaaccttggttcatagggtgttt  
tcatgaactttatgttcatgtgtattttactattattagaggtctatcttcagagaggag  
tacaagaaatgggattactgggtgcaaaggtaaatggatatgtgtctttgctaggtattg  
ccaaattttatctccagaaatcttgcacaaatctgtactcctgttagcaatgtgtgctgat  
acctgcttaccacatgacctcagtaaaagaatgtgttgtcatattgggtattgaaatttta  
gcactgtaagcaacagggtcattttggaaaacctgagctttcgccaaattcagctattttg  
atttgctttttattattagcatataccaaaaataaataggcatattagagtttctttcttg  
catcttaaaattcatctaacacatctataataacattcttttctttttttccattctag

GACTTGCCCCCTTTCGTCTATTTGTCAGACGAATGTTACAATTTACTGGCAATAAAGTTTT  
L A P F V Y L S D E C Y N L L A I K F W 3106

GGATAGACCTTAATGAGGACATTATTAAGCCTCATATGTTAATTGCTGCAAGCAACCTCC  
I D L N E D I I K P H M L I A A S N L Q 3126

AGTGCGCACCAGAATCCAAATCAGGCCTTCTTACTTTATTTGCTGGAGATTTTCTGTGT  
W R P E S K S G L L T L F A G D F S V F 3146

TTTCTGCTAGTCCAAAAGAGGGCCACTTTCAAGAGACATTCAACAAAATGAAAAATACTG  
S A S P K E G H F Q E T F N K M K N T V 3166

TTGAGgtaagggttacttttcagcatcaccacacattttgggtatttttctattttgacagt  
E 3167

SUBSTITUTE SHEET (RULE 26)

# Fig.7. (20/22) <sup>37/52</sup>

ccagtatcaaggaaatagctttttatacaaaattggatagttgaggtagttatgtgaggtaaa  
gtttaatcatatattaattgcccataaacctcaggagatgggggaatggggaaatgacag  
cagctagaaagagaagaatgacttgaagggaatgagttaggagaaattgtgagaaggat  
gttcagaaatgcagactttgtaagcaaaactggaaattgggttacaagaataatagagtt  
tctgtggtttgcagcagtcagcagtggtgattagtttaataatagagactacaggtttac  
atttaaactccatattctagtggttttatacagattatatttctttgacttgatttaatccc  
agataagagacactgatattattttccctagatcatgt.....

## Exon 26

.....tatttgaccagttgtgattt  
ctcaagcaacagtttcatgtagtttaacctataaatcatttcaattlaattcttgyaacag  
acgtgaggttaggtgaggcaattctttcttttcttaaccaaagaagtacctttatagatg  
tgagatgattcccagctattaagtagtaaatagagctaggacttgagccccaatcttcca  
gcttcaatccagatcatatgacagcttgctgattaaactagatgacagagaagatctctt  
tccttcagatacacatactttttctctgttccccctctccctatcagctagattcccctaa  
atcactgatactgggttttgaattttgcatcggcatgtttgacaattgggtatcacattta  
gggtttttcattcttttttgggtcccaaacttttcatttctgcttttaaggaaatacttt  
tggaaacataaatatgtgggtttgcaattttataaagcagcttttccacttattttcttag

AAATATTGACATACTTTGCAATGAAGCAGAAAAACAAGCTTATGCATATACTGCATGCAAAT

N I D I L C N E A E N K L M H I L H A N 3187

GATCCCAAGTGGTCCACCCCACTAAAGACTGTACTTCAGGGCCGTACACTGCTCAAATC

D P K W S T P T K D C T S G P Y T A Q I 3207

ATTCCCTGGTACAGGAAACAAGCTTCTGgtaagttaatgtaaactcaaggaatattataag

I P G T G N K L L 3216

aagtatatatggaggccatcgatatattctgttgatatacctagtaaacatgggtaaaatgta  
attaaacttaattagaaaatgtgggttggtatgtggctcctgtaagtatagttatttagaa  
attttatttattgaagcaagatatgaaactctgggtgcacactttccaaacaggtgcttt  
catttacatgtgattgaaaagtgttttttgccatttatttcaactgttccatacaattagg  
gttggtttctaagctgtttgtaagctgtttctaagctattttaagtggttaaatcacagtag  
atgcaaatcaagctaaaagtctttaacattggctaatggctgattctttaaatagctaaatc  
ttgctaagggtatctatatattaactcatttaactcctcataacaaccctatgagataaaacc  
taagtcctcacttaacattgtcaataggttttttgaaactgatttta.....

## Exon 27

taatctggt  
aataataaaaaaacaagattaaagcataagtgacgtccccctacctccttttttatcttt  
tactgtgattattcttcatcttcttcttcttcttcttcatgtcattttatatgttctttagtaa  
attactttcatctagaataggaataatgtgaaactgaaatcacctaacctattaggagttta  
ggggaggaggactgtgtgtaatatattgctgtcttaaatattttcaatgaaaagttacttt  
gatttagttttttatgttactacataattatgataggctacgtttttcatttttttatcag

ATGTCTTCTCCTAATTGTGAGATATATTATCAAAGTCCTTTATCACTTTGTATGGCCAAA

M S S P N C E I Y Y Q S P L S L C M A K 3236

AGGAAGTCTGTTTCCACACCTGTCTCAGCCCAGATGACTTCAAAGTCTTGTAAAGGGGAG

R K S V S T P V S A Q M T S K S C K G E 3256

Fig.7. (21/22)

38/52

AAAGAGATTGATGACCAAAGAAGCTGCAAAAAGAGAAGAGCCTTGGATTTCTTGAGTAGA  
K E I D D Q K N C K K R R A L D F L S R 3276

CTGCCTTTACCTCCACCTGTTAGTCCCATTGTGACATTTGTTTCTCCGGCTGCACAGAAG  
L P L P P P V S P I C T F V S P A A Q K 3296

GCATTTTCAGCCACCAAGGAGTTGTGGCACCAAATACGAAACACCCATAAAGAAAAAAGAA  
A F Q P P R S C G T K Y E T P I K K K E 3316

CTGAATTCTCCTCAGATGACTCCATTTAAAAAATTCAATGAAATTTCTCTTTTGGAAAGT  
L N S P Q M T P F K K F N E I S L L E S 3336

AATTCAATAGCTGACGAAGAAGCTTGCAATTGATAAATACCCAAGCTCTTTTGTCTGGTTCA  
N S I A D E E L A L I N T Q A L L S G S 3356

ACAGGAGAAAAACAATTTATATCTGTCTAGTGAATCCACTAGGACTGCTCCCACCAAGTTCA  
T G E K Q F I S V S E S T R T A P T S S 3376

GAAGATTATCTCAGACTGAAACGACGTTGTACTACATCTCTGATCAAAGAACAGGAGAGT  
E D Y L R L K R R C T T S L I K E Q E S 3396

TCCCAGGCCAGTACGGAAGAATGTGAGAAAAATAAGCAGGACACAATTACAATAAAAAA  
S Q A S T E E C E K N K Q D T I T T K K 3416

TATATCTAAGCATTGTGCAAAGGCGACAATAAATTATTGACGCTTAACCTTTCCAGTTTAT  
Y I STOP 3418

AAGACTGGAATATAATTTCAAACCACACATTAGTACTTATGTTGCACAATGAGCAAAGAA  
ATTAGTTTCAAATTTACCTCAGCGTTTGTGTATCGGGCAAAAATCGTTTTGCCCGATTCC  
GTATTGGTATACTTTTGCTTCAGTTGCATATCTTAAACTAAATGTAATTTATTAATAA  
TCAAGAAAAACATCTTTGGCTGAGCTCGGTGGCTCATGCCTGTAATCCCAACACTTTGAG  
AAGCTGAGGTGGGAGGAGTGCTTGAGGCCAGGAGTTCAAGACCAGCCTGGGCAACATAGG  
GAGACCCCCATCTTTACAAAGAAAAAAGGGGAAAAAGAAAATCTTTTAAATCTTTGG  
ATTTGATCACTACAAGTATTATTTTACAAGTGAATAAACATACCATTTTCTTTTAGATT  
GTGTCATTAAATGGAATGAGGTCTCTTAGTACAGTTATTTTGATGCAGATAATTCCTTTT  
AGTTTAGCTACTATTTTAGGGGATTTTTTTTAGAGGTAACCTCACTATGAAATAGTTCTCC  
TTAATGCAAATATGTTGGTTCTGCTATAGTTCCATCCTGTTCAAAAGTCAGGATGAATAT  
GAAGAGTGTTGTTTCCTTTTGAGCAATTCTTCATCCTTAAGTCAGCATGATTATAAGAAA  
AATAGAACCCTCAGTGTAACCTCTAATTCCTTTTACTATTCCAGTGTGATCTCTGAAATT  
AAATTACTTCAACTAAAAATTCAAATACTTTAAATCAGAAGATTTTCATAGTTAATTTATT

39/52

Fig.7. (22/22)

TTTTTTTTcaacaaaatgggtcatccaaactcaaaacttgagaaaatatcttgcctttcaaat

tggcactgattctgcctgctttatTTTTtagcgctatcacaggacccagagcctatgcctt  
tttaaacttaccacaaaagcagaagattaattcaatttaagatgatactctcatttggtta  
cgtccctTTTTTTTTTTTTTggagatggagcttgcctttgtcgcccatgctggagtgag  
tggcatgatccctggctcactgcagcctccacttcccgggttcaagtaattctcccacctc  
aagcctccctagtagctgggattacagggagcgcaccaccatgccagctaatttttgcatt  
tttttagtagagactgggttttaccatggttgccaagctgggtctcaaaactcctgatgtcag  
gtgatccatctgcctcagcctcccaaagtgcctgggttatagggctgagccactgtgccc  
ggccaatatTTTgttactttcttaggtttaatagagaaaagggataaaaacatttctaactg  
ggagttaatagcatggagaaggtcttaaatcagatggttttaatgccttaaatgtctgtat  
aatatcatgTTTTcaaatctaattataaaatcgtttaaagccaagaataaatctTTTTaa  
aaattgacttgTTTTccttccataactctgagccatgatttttctgttctgtaaaaagcat  
taacaaaattgtctatTTTgttactccctgtaacttaagtattctgcaagctcttattaat  
gagacttgTTTTgtttctaaaacagtttggttttcacatcctaattttgcagtgatccac  
tctagaacaaggaaaataaaacttggttttcaaacaggagaaacagaaaaaattacaagaat  
ttaacctTTTTTTTTTgaatccttggtacaactgctattgtctgttctcatgtagaaca  
ccattatgttgatagatatgtaatgcgcacactTTTTtataaattataaataacatcca  
agctacatgaaacaaaatatgaagcttgagtatatgtgcataattatccccctcaaaagtga  
caatttaatgactacaaagtcaacatactgcatacataagggataatggtaaaaatttggtg  
ttatttatcagcaatcatttaatgagaccctattgcccataaagagcatttgccagttc  
tatgaatgatgcaaacagcgcaacacaatacaagtcataattgggtgttcaaagagttaca  
caaatcagaagccatgggagttgaaaaacagatcacttctacaggggataaaagtgtgatgg  
catttaatcaattTTTTatgggttacaaaccaaataatttcaggggtcagctagggaaagaa  
gagacattaaataggccaggaaactattcttttagagctctagaaaagttatgtaaccag  
agccacttctccatagtggtcactcacttaaccecaagtgcagcca.....

40/52

Fig.8. (1/4)

## BRCA2 SSCP PRIMERS

EXON	PRIMER F	PRIMER R	SIZE (bp)	CON
1				
2	CTCAGTCACATAATAAGGAAT	ACACTGTGACGTACTGGGTTTT	256	TD60-50
3	TCCTGGGTCACAAAATTGCTGTCA	TTCCCTAGTTTGTAGTTCTCCCCAGTC	356	TD60-50
4	AGAATGCAAAATTTATAATCCAGAGTA	AATCAGATTTCATCTTTATAGAACAAA	249	TD60-50
5	AACAATTATATGAATGAGAATC	AATTGTTAAGTTTATTATTTTATTA	220	TD55-45
6	CCACAAAGAGATAAGTCAGGTA	TGTAATCTCAGGGCAAAAGGTA	234	TD55-45
7	TAACTGAAATAAAGAGTGAA	AACAGAAGTATTAGAGATGAC	275	TD55-45
8	AATAGTAGATGTGCTTTTIGA	ACATATAGGACCAGGTTTAGAGAC	285	TD60-50
9	GTTCAACTAAACAGAGGACT	CTAGTGATTTTAAACTATAATTTTGG	164	TD68-60
10A	TATAAATATTAATGTGCTTCTGTT	AAAGGGCTTCTGATTTGCTAC	374	TD55-45

4/52

Fig.8. (2/4)

10B	ATCTGAAGTGAACCAATGATAC	ACGTGGCAAGAATTCCTGAAGTAA	270	TD60-50
10C	TTTCAGAAAAAGACCTATTAGACA	CTTTTGTGATACCTGAAATGAAGAAG	260	TD60-50
10D	TAAAGCAGGCAATATCTGGAACCTCT	GTGGATATTAAACCTGCATTCTTCAA	280	TD60-50
10E	TATGTCCAAATTTAATTGATAAT	AAACACAGAAGGAATCGTCATC	320	TD55-45
11A	GATGGTACTTTAAATTTTGTCACTTTG	TTCCCTTATTACATTTTGTCTCTTTAT	220	TD55-45
11B	CTAGCTCTTTTGGGACAATTTCTGAGG	TTTCATGATCATATAAAGANT	300	TD55-45
11C	CAAAAGTGAATACAGTGATAC	ATAATTTTCATTTAAAGCACATACAT	250	TD60-50
11D	TCTAGAGGCAAGAATCATTA	TCATTGCTTGAGAAAAGTTC	305	TD55-45
11E	TTCAAAAATAACTGTCAATCC	GTGCTTGTTTTATCACCTGT	230	TD60-50
11F	AACCCATTTTCAAGAACTCTACCA	CTGAAGCTACCTCCAAAACCTGTG	274	TD60-50
11G	ACAAATGGCAGGACTCTTAGG	CCTGCTTGGAAAAATAACATCTG	300	TD60-50
11H	AGTTGTTTCTGATTTGPAAAAATAGTC	GCTGCTGTCTACCTGACCAA	310	TD60-50
11I	GATGCTGATCTTCATGTCATAA	TACTCTGCAGAAGTTTCCTCACCTA	300	TD60-50
11J	GGGTTTAGGGCTTTTATTTC	TATCAGTTGGCATTATTATTATTTT	237	TD60-50
11K	CTTCAAGTAAATGTCATGATCTGTC	CTGGCAGCAGTATATTGTTATCT	202	TD60-50
11L	ATTACTGMAAATTACAAGAGAAATA	AAAAAGTTAAATCTGACTCAAAATCT	244	TD60-50
11M	TTTATGAAGGAGGGAACACTCA	TTCAAGAAATTAAGGAAAAGTTATGC	282	TD60-50
11N	AAAAATATTAGTGTGCCCAAAGAG	TGATGAAAACCCACACAGATAGG	278	TD60-50
11O	GTAAACACACAAAATACTGAAAG	CATTGATGGCTAAAACCTGGTG	233	TD60-50
11P	TCATACAGCTAGCGGAAAAA	TGGCACCAACAGTCTCAATAG	265	TD60-50
11Q	CTGCCCCAAAGTCTAAAGAAAT	AATGACTGAATAAGGGGACTGAT	230	TD60-50
11R	GTCCCTGCAACTTGTTCAC	GATTTTGTTCATTTTCAGC	230	TD60-50
11S	AACCAGAAAGATAAATACT	TCCTCAACGCAAAATATCTTCAT	334	TD55-45
11T	TTTCCAAAAGTAATATCCAAATGTA	ATTTTGTATTATCTCTCGTTGTT	291	TD55-45
11U	AAGACATATTACAGACAGT	TGAAGCTTCCCTATACATACAT	300	TD60-50
11V	CACCTTGTGATGTTAGTTTG	TTGGGATATTAAATGTTCTGGAGTA	350	TD60-50
11W	AAAGTAACGAACATTCAGACCA	CTGGGTTTCTCTTATCAACNCGA	310	TD60-50
11X	AGCTTTCACATATTCACCTACG	GTGAGACTTTGGTTCCCTAAT	280	TD60-50
11Y	TTCAACAAGACAAACACAGT	TGTCAGTTTCATCATCTTCCATAAA	270	TD60-50
11Z	TTACTCCAAAGATTCAGAAAACCTAC	AGCATACCAAGTCTACTGAATAAAC	290	TD60-50



42/52

Fig.8. (3/4)

12	AGGTCACATATTGTTGTAAG	AGTGGCTCATGCTGTGAAT	358	TD60-50
13	TAAAGCCTATAAATTGCTCA	CTTCTTAACGTTAGTGCAAT	271	TD55-45
14A	ATGTAGCAAAATGAGGGTCTG	GTCTGCCTGTAGTAATCAAGTGT	289	TD60-50
14B	CTTCAAGCAATTTAGCAGTTTCAG	TGCTGCTTGATTTGGAGTTGTT	292	TD55-45
14C	TATTAACTTGGAGGAAAACAGACA	CAAAGGGGAAAACCATCAG	250	TD60-50
15	GGCCAGGGGTTGTGCTTTT	AGGATACTAGTTAATGAATA	314	TD60-50
16	TTTGGTAAATTCAGTTTGGTTT	AACACACAATCTTTTTCATAGA	330	TD60-50
17	CAGAGAATAGTTGTAGTTGTTGAA	AGAAACCTTAACCCCACTATGC	306	TD60-50
18A	ATTTCAGTTTATTTCTCAGTTATTC	ACCACCCATCTGTAAGTTCA	296	TD60-50
18B	ATCTGAAACTTCTAGCAATAA	TTTAACTGAATCAATGACTG	281	TD60-50
19	AAGTGAATATTTTAAAGGCAGTT	TATATGGTAAGTTTCAAGAAT	296	TD60-50
20	CACGTGTCCTGGCCTGATAC	ATGTTAAATTCAAAGTCTCTA	296	TD60-50
21	GGGTGTTTATGCTTGGTTCT	CATTTCAACATATTTCCCTTCCTG	304	TD60-50
22	TTTTGTTCTGATGCTTTTATTC	AATCATTTTGTAGTAAGTTCAT	314	TD60-50
23	ACTTCTTCCATTCATCTTTCTCA	AAAACAAAACAAAAATTC AACATA	290	TD60-50
24	CAGTTTGTGATAAGTCTTGT	AGCTCCAACATAATCAFAAGA	290	TD60-50
25A	TAAATTCATCTAACACATCTAT	ACCTCAACAGTATTTTTCATTT	302	TD55-45
25B	AATCCCAAAATCAGGCCTTCTTACTT	ATTTCCCCATTTCCCCCATCT	283	TD60-50

43/52

Fig.8. (4/4)

26	AAGGAATACCTTTGGAAACATAA	TTTACTAGGTATACACAGAA	299	TD55-45
27A	TAGGAGTTAGGGAGGGAGACTGTGT	TTTCTCCCTTTTACAAGACTTTGAAG	252	TD60-50
27B	TTGTATGGCCAAAGGAAGTCTG	TTTTTAAATGGAGTCATCTGAGGAGA	285	TD60-50
27C	AAGGCAITTCAGCCACCACCAAGGAGT	CAACGTCGTTTCAGTCTGAGATAATC	272	TD60-50
27D	AGGAGAAAAACAATTTATATCTGICA	GTGGTTTGAAATTATATTCAGTCTT	264	TD60-50

Fig.9.

## BRCA2 PTT primers

Exon	Primers	PTT primer sequence	Size
10	JAD20F CAAAGACCACATTGGAAAGTC x JAD20R GCAAAATGTAAGTGTGCTTC	PTT12 GAAACAGTTGTACATACCTCTGAAGA	1kb approx. 37kd
11	JAD53F GATGGTACTTTTAA'TTTT'GTAC x JAD13R AAATGACTCTTTTGGCGACAC	PTTD2 GA'TTCTGAAAGAACCAACTTTGTCC	2.4kb approx. 89kd
11	JAD12F AGATTTTGAGACTTCTGATAC x JAD1R AGCATACCAAGTCTACTGAATAAAC	PTTA GACATTCTAAAGTTATGAGGA	2.4kb approx. 89kd
27	JAD43F ACATAATTATGATAGGCTACG x JAD45R AAGCGTCAATAATTTATTGTC	PTTJ TCTTCTCCTAATTGTGAGATA	0.6kb approx. 22kd

44/52



Fig.10. (2/4)

INTVSHLQSSVVVSDCEISHATQMLSSKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	430	440	450	460	470	480	490	500	510	520	530	540	550	
INTVSHLQSSVVVSDCNSHITQMLFSKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	511													
INTVSHLQSSVVVSDCNSHITQMLFSKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	503													
INLISGFVQNSTVSDSESHATPPTLSLKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	543													
VTMSHPQSQSVS - CEDTHISLQVLPKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	3													
VTVTSQSHNQSVS - HEDITDIAQMLSSKQOFNSNINLTPSOKAEITELSTILESSQOFETQFRKPESHIAQKNITFEPENQMTILNTTSEBMDADLHLTVNAPSI	484													
MOUSE . PRO	528													
QVDSKQFBSAGIKQKFACLLKSSONKASGYLTDENEVEFRGFYSALGTIKLVSSPALQKAVKLFSDIENISETSAMDPIISLSSKCHDSVSMFKIENQN-DKS	560	570	580	590	600	610	620	630	640	650	660	670	680	
HUMAN . PRO	620													
MONKEY . PRO	612													
DOG . PRO	649													
PIG . PRO	29													
HAMSTER . PRO	594													
MOUSE . PRO	637													
SEKNNKQQLILQNNIEMITGIFVEENTENKENTENEDNKYTCASRNS - NLE - SDGSSSKNDTVYIHKQETDLPFDQ - NICLKLSQOFMBGNTQIKEDLSDLTFL	690	700	710	720	730	740	750	760	770	780	790	800	810	
HUMAN . PRO	728													
MONKEY . PRO	719													
DOG . PRO	757													
PIG . PRO	135													
HAMSTER . PRO	704													
MOUSE . PRO	742													
VMAEETCHGVNSKEQJATATKTEQNIKDFDIFDISQFASGRNIRVSKESLNKAVNFFDQ - TTEELNFSDSLSELGNNKNDISSHE - - - ETDIVNKNILKES	820	830	840	850	860	870	880	890	900	910	920	930	940	
HUMAN . PRO	832													
MONKEY . PRO	823													
DOG . PRO	866													
PIG . PRO	242													
HAMSTER . PRO	806													
MOUSE . PRO	843													


47/52

**SUBSTITUTE SHEET (RULE 26)**

Fig. 10. (4/4)

NSLDSBCSTNSHKVFADIQSEIILQHNQSMGIEKVEIPPOVSLSTSDICKSIGKLPKSVSSNTGCIFFSTASGSKVWSDASIQARQVFSIEDSAKQLFSKV	1330	1340	1350	1360	1370	1380	1390	1400	1410	1420	1430
NSLNDCESTHSHKVFADIQSEIILQHNQSMGIEKVEIPPOVSLSTSDICKSIGKLPKSVSSNTGCIFFSTASGSKVWSDASIQARQVFSIEDSAKQLFSKV	1335										
NSLDSBCSMHSHKVFADIQSEIILQHNQSMGIEKVEIPPOVSLSTSDICKSIGKLPKSVSSNTGCIFFSTASGSKVWSDASIQARQVFSIEDSAKQLFSKV	1340										
NSLGAIE-CSPSHKVFADIQSEIILQHNQSMGIEKVEIPPOVSLSTSDICKSIGKLPKSVSSNTGCIFFSTASGSKVWSDASIQARQVFSIEDSAKQLFSKV	1345										
NSPSEEHITRSQVFPFIQSEIILQHNQSMGIEKVEIPPOVSLSTSDICKSIGKLPKSVSSNTGCIFFSTASGSKVWSDASIQARQVFSIEDSAKQLFSKV	1350										
AL--RDD-YMNSHKILFTHDEQIILQHNQSMGIEK--AQIPP--VLEIWDTSK-STRELAQAACSSHMGIFSTASGKAVVSDASLEKARQVFSIEDSAKQLFSKV	1355										
SS--SEJVCINSRKDSFCHNEQIILQHNQSMGIEK--AQIPP--VLEIWDTSK-STRELAQAACSSHMGIFSTASGKAVVSDASLEKARQVFSIEDSAKQLFSKV	1360										
LAKNE-HSDQLTREENTVHTPENILLSSQK-FS-NVWSSAFSGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1440	1450	1460	1470	1480	1490	1500	1510	1520	1530	1540
FKSNE-HSDQLTREENTVHTPENILLSSQK-FS-NVWSSAFSGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1445										
FKSNE-HSDQLTREENTVHTPENILLSSQK-FS-NVWSSAFSGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1450										
LAKNEHSEKFTNEENTVIYTSQNLSS-----AFSGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1455										
VKSDEHSDKYTREENTVHTPENILLSSQK-FS-NVWSSAFSGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1460										
LESHE-QSDHSGRRNSVTHNPEDVLSLPKTPA-SVANSVFSFGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1465										
LEBNE-KPHISVARENSVWHSQTQGVLSLPKPLP-QVWSSVFSFGFSTASGKQVSVSESALHKVKGMLEEFDLIRTEISLQHSPTSRQDVSKILFPQSCVDKRTPEHSVNS	1470										
LEKTYDEFSLENNVWSSGSENNHISIKVSPQLSQFKQKQ-QLVLGTRKVSLENNHIVLCKEQTSPENWMEIGKTETETFSVPVKTIN-----YSK-----	1550	1560	1570	1580	1590	1600	1610	1620	1630	1640	1650
MEKTCSEKFTLSNNLWBGSSSENNHISIKVSPQLSQFKQKQ-QLVLGTRKVSLENNHIVLCKEQTSPENWMEIGKTETETFSVPVKTIN-----YSK-----	1555										
MEKTCSEKFTLSNNLWBGSSSENNHISIKVSPQLSQFKQKQ-QLVLGTRKVSLENNHIVLCKEQTSPENWMEIGKTETETFSVPVKTIN-----YSK-----	1560										
MEKTCSEKFTLSNNLWBGSSSENNHISIKVSPQLSQFKQKQ-QLVLGTRKVSLENNHIVLCKEQTSPENWMEIGKTETETFSVPVKTIN-----YSK-----	1565										
TEKAYNBQFKLPDSNTSESSSENNHISIKVSPQLSQFKQKQ-QLVLGTRKVSLENNHIVLCKEQTSPENWMEIGKTETETFSVPVKTIN-----YSK-----	1570										
LQKTYDOKFSLPNNYK-ESASLGNTSHLEASFOI SQF--KQDTRMLGTRKVS-----LLEKQJFTFQNTKTESQMET	1575										
LQKTYDOKFSLPNNYK-ESASLGNTSHLEASFOI SQF--KQDTRMLGTRKVS-----LLEKQJFTFQNTKTESQMET	1580										
LQKTYDOKFSLPNNYK-ESASLGNTSHLEASFOI SQF--KQDTRMLGTRKVS-----LLEKQJFTFQNTKTESQMET	1585										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1590										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1595										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1600										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1605										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1610										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1615										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1620										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1625										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1630										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1635										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1640										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1645										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1650										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1655										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1660										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1665										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1670										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1675										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1680										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1685										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1690										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1695										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1700										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1705										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1710										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1715										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1720										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1725										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1730										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1735										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1740										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1745										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1750										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1755										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1760										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1765										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1770										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1775										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1780										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1785										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1790										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1795										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1800										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1805										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1810										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1815										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1820										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1825										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1830										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1835										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1840										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1845										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1850										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1855										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1860										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1865										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1870										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1875										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1880										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1885										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1890										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1895										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1900										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1905										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1910										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1915										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1920										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1925										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1930										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1935										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1940										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1945										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1950										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1955										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1960										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1965										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1970										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1975										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1980										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1985										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1990										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	1995										
AKAFMEDELTDSE-QTHAKCSLFTCTQNETLFSNRTRKRRTMTVDV	2000										


Fig.11.






59/52

Fig.12.



5/52

Fig.13A.



# INTERNATIONAL SEARCH REPORT

Internat'l Application No  
PCT/GB 96/02904

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>GENOMICS, 1996, 36, 86-99, XP000647994            COUCH FJ ET AL: "GENERATION OF AN            INTEGRATED TRANSCRIPTION MAP OF THE BRCA2            REGION ON CHROMOSOME 13Q12-Q13"            see the whole document            -----</p>	<p>1-10,            14-29</p>


## Information on patent family members

PCT/GB 96/02904

Form PCT/ISA/210 (patent family annex) (July 1992)

52/52

Fig.13B.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 96/02904

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K14/82 C12Q1/68 G01N33/53 A61K38/17 A61K48/00  
A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SCIENCE, SEP 30 1994, 265 (5181) P2088-90, UNITED STATES, XP002029509 WOOSTER R ET AL: "Localization of a breast cancer susceptibility gene, BRCA2, to chromosome 13q12-13."  see the whole document ---	1-10, 14-37, 39-46, 52-56, 60-62, 64-66
Y	SCIENCE, OCT 7 1994, 266 (5182) P66-71, UNITED STATES, XP000202410 MIKI Y ET AL: "A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1."  see the whole document ---	1-10, 14-37, 39-46, 52-56, 60-62, 64-66
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

15 April 1997

Date of mailing of the international search report

25. 04. 97

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

# INTERNATIONAL SEARCH REPORT

Intern. al Application No  
PCT/GB 96/02904

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95 19369 A (UNIV VANDERBILT) 20 July 1995  see the whole document ---	1-10, 14-37, 39-46, 52-56, 60-62, 64-66
A	CANCER RES, NOV 1 1995, 55 (21) P4830-2, UNITED STATES, XP002029510 GUDMUNDSSON J ET AL: "Different tumor types from BRCA2 carriers show wild-type chromosome deletions on 13q12-q13." see the whole document ---	1-10, 14-33
A	PROC NATL ACAD SCI U S A, JUN 20 1995, 92 (13) P5950-4, UNITED STATES, XP002029511 SCHUTTE M ET AL: "Identification by representational difference analysis of a homozygous deletion in pancreatic carcinoma that lies within the BRCA2 region." see the whole document ---	1-10, 14-33
P,X	NATURE, 1995, 378, 789-792, XP002029512 WOOSTER R ET AL: "IDENTIFICATION OF THE BREAST-CANCER SUSCEPTIBILITY GENE BRCA2" see the whole document ---	1-10, 14-29
P,X	CANCER RESEARCH, 1996, 56, 3622-3625, XP002029513 FOSTER KA ET AL: "SOMATIC AND GERMLINE MUTATIONS OF THE BRCA2 GENE IN SPORADIC OVARIAN-CANCER" see the whole document ---	1-10, 14-29
P,X	CANCER RESEARCH, 1996, 56, 3409-3414, XP002029514 CORIXA CORPORATION: "A COMMON MUTATION IN BRCA2 THAT PREDISPOSES TO A VARIETY OF CANCERS IS FOUND IN BOTH JEWISH ASHKENAZI AND NON-JEWISH INDIVIDUALS" see the whole document ---	1-10, 14-29
P,X	ONCOGENE, 1996, 13, 213-221, XP000618318 JACOB ANK ET AL: "ISOLATION OF EXPRESSED SEQUENCES THAT INCLUDE A GENE FOR FAMILIAL BREAST-CANCER (BRCA2) AND OTHER NOVEL TRANSCRIPTS FROM A 5-MEGABASE REGION ON CHROMOSOME 13Q12" see the whole document ---	1-10, 14-29
	---	

-/--